

β-catenin (BDI710): sc-59894

BACKGROUND

The catenins, α , β and γ , are proteins which bind to the highly conserved, intracellular cytoplasmic tail of E-cadherin. Together, the catenin/cadherin complexes play an important role mediating cellular adhesion. α -catenin was initially described as an E-cadherin associated protein, and since has been shown to associate with other members of the cadherin family, such as N-cadherin and P-cadherin. β -catenin associates with the cytoplasmic portion of E-cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. β -catenin has also been found in complexes with the tumor suppressor protein APC. γ -catenin, also known as plakoglobin, binds with α -catenin and N-cadherin. It has been shown that the transmembrane phosphatase PTP μ associates with catenin/cadherin complexes and may regulate complex signaling.

REFERENCES

- Knudsen, K.A., Soler, A.P., Johnson, K.R. and Wheelock, M.J. 1995. Interaction of α -Actinin with the cadherin/catenin cell-cell adhesion complex via α -catenin. *J. Cell Biol.* 130: 67-77.
- Breen, E., Steele, G. Jr. and Mercurio, A.M. 1995. Role of the E-cadherin/ α -catenin complex in modulating cell-cell and cell-matrix adhesive properties of invasive colon carcinoma cells. *Ann. Surg. Oncol.* 2: 378-385.
- Pierceall, W.E., Woodard, A.S., Morrow, J.S., Rimm, D. and Fearon, E.R. 1995. Frequent alterations in E-cadherin and α - and β -catenin expression in human breast cancer cell lines. *Oncogene* 11: 1319-1326.
- Takayama, T., Shiozaki, H., Shibamoto, S., Oka, H., Kimura, Y., Tamura, S., Inoue, M., Monden, T., Ito, F. and Monden, M. 1996. β -catenin expression in human cancers. *Amer. J. Pathol.* 148: 39-46.
- Ozawa, M., Nuruki, K., Toyoyama, H. and Ohi, Y. 1995. Cloning of an alternative form of plakoglobin (γ -catenin) lacking the fourth armadillo repeat. *J. Biochem.* 118: 836-840.
- Sacco, P.A., McGranahan, T.M., Wheelock, M.J. and Johnson, K.R. 1995. Identification of plakoglobin domains required for association with N-cadherin and α -catenin. *J. Biol. Chem.* 270: 20201-20206.
- Brady-Kalnay, S.M., Rimm, D.L. and Tonks, N.K. 1995. Receptor protein tyrosine phosphatase PTP μ associates with cadherins and catenins *in vivo*. *J. Cell Biol.* 130: 977-986.

CHROMOSOMAL LOCATION

Genetic locus: CTNNB1 (human) mapping to 3p21; Ctnnb1 (mouse) mapping to 9 F4.

SOURCE

β -catenin (BDI710) is a mouse monoclonal antibody raised against full length β -catenin of human origin.

PRODUCT

Each vial contains 50 μ g IgG_{2a} in 500 μ l of PBS with < 0.1% sodium azide, 1% gelatin, PEG and sucrose.

APPLICATIONS

β -catenin (BDI710) is recommended for detection of β -catenin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 μ g per 100–500 μ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for β -catenin siRNA (h): sc-29209, β -catenin siRNA (m): sc-29210, β -catenin siRNA (h2): sc-44252, and β -catenin siRNA (m2): sc-44253.


Molecular Weight of β -catenin: 92 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, A-431 whole cell lysate: sc-2201 or MCF7 whole cell lysate: sc-2206.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



β -catenin (BDI710): sc-59894. Western blot analysis of β -catenin expression in non-transfected: sc-117752 (A) and human β -catenin transfected: sc-116622 (B) 293T whole cell lysates.



β -catenin (BDI710): sc-59894. Western blot analysis of β -catenin expression in MCF7 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.