

Integrin α IIb (54B3): sc-59967

BACKGROUND

Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. The 16 α and 8 β subunits heterodimerize to produce more than 20 different receptors. Most integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, Collagen and Vitronectin. Certain integrins can also bind to soluble ligands such as Fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis.

REFERENCES

- Hynes, R.O. 1992. Integrins: versatility, modulation and signaling in cell adhesion. *Cell* 69: 11-25.
- Miyamoto, S., Akiyama, S.K. and Yamada, K.M. 1995. Synergistic roles for receptor occupancy and aggregation in integrin transmembrane function. *Science* 267: 883-885.
- Clark, E.A. and Brugge, J.S. 1995. Integrins and signal transduction pathways: the road taken. *Science* 268: 233-239.
- Sheppard, D. 1996. Epithelial integrins. *Bioessays* 18: 655-660.
- Juliano, R. 1996. Cooperation between soluble factors and integrin-mediated cell anchorage in the control of cell growth and differentiation. *Bioessays* 18: 911-917.
- Hantgan, R.R., Lyles, D.S., Mallett, T.C., Rocco, M., Nagaswami, C. and Weisel, J.W. 2003. Ligand binding promotes the entropy-driven oligomerization of Integrin α IIb/ β 3. *J. Biol. Chem.* 278: 3417-3426.
- Goncalves, I., Hughan, S.C., Schoenwaelder, S.M., Yap, C.L., Yuan, Y. and Jackson, S.P. 2003. Integrin α IIb/ β 3-dependent calcium signals regulate platelet-fibrinogen interactions under flow. Involvement of phospholipase C γ 2. *J. Biol. Chem.* 278: 34812-34822.
- Maxwell, M.J., Yuan, Y., Anderson, K.E., Hibbs, M.L., Salem, H.H. and Jackson, S.P. 2004. SHIP1 and Lyn kinase negatively regulate Integrin α IIb/ β 3 signaling in platelets. *J. Biol. Chem.* 279: 32196-32204.
- Huang, C.L., Cheng, J.C., Liao, C.H., Stern, A., Hsieh, J.T., Wang, C.H., Hsu, H.L. and Tseng, C.P. 2004. Disabled-2 is a negative regulator of Integrin α IIb/ β 3-mediated fibrinogen adhesion and cell signaling. *J. Biol. Chem.* 279: 42279-42289.

SOURCE

Integrin α IIb (54B3) is a mouse monoclonal antibody raised against a 32 amino acid synthetic peptide corresponding to the cytoplasmic domain of Integrin α IIb of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Integrin α IIb (54B3) is recommended for detection of the cytoplasmic domain of Integrin α IIb of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Integrin α III siRNA (h): sc-63319, Integrin α III siRNA (m): sc-63320, Integrin α III shRNA Plasmid (h): sc-63319-SH, Integrin α III shRNA Plasmid (m): sc-63320-SH, Integrin α III shRNA (h) Lentiviral Particles: sc-63319-V and Integrin α III shRNA (m) Lentiviral Particles: sc-63320-V.

Molecular Weight of Integrin α IIb: 118 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.