

AP-3 μ siRNA (h): sc-60177

BACKGROUND

Clathrin-coated pits and vesicles are assembled for receptor-mediated endocytosis through interaction with Clathrin associated protein complexes. Vesicle transport is mediated from the *trans*-Golgi network by the adapter complex AP-1 and from the plasma membrane by the AP-2 complex. AP-3 (also designated AP180 or F1-20) is a synapse-specific Clathrin assembly protein. The protein CALM (Clathrin assembly protein lymphoid myeloid leukemia) is highly homologous to AP180 and may also be involved in Clathrin assembly. AP-3 δ , AP-3 σ and AP-3 μ are important parts of the AP-3 complex.

REFERENCES

1. Robinson, M.S. 1989. Cloning of cDNAs encoding two related 100-kD coated vesicle proteins (α -Adaptins). *J. Cell Biol.* 108: 833-842.
2. Kirchhausen, T., et al. 1989. Structural and functional division into two domains of the large (100- to 115-kDa) chains of the Clathrin-associated protein complex AP-2. *Proc. Natl. Acad. Sci. USA* 86: 2612-2616.
3. Robinson, M.S. 1990. Cloning and expression of γ -Adaptin, a component of Clathrin-coated vesicles associated with the Golgi apparatus. *J. Cell Biol.* 111: 2319-2326.
4. Ponnambalam, S., et al. 1990. Conservation and diversity in families of coated vesicle adaptins. *J. Biol. Chem.* 265: 4814-4820.
5. Simpson, F., et al. 1997. Characterization of the adaptor-related protein complex, AP-3. *J. Cell Biol.* 137: 835-845.
6. Singh, B., et al. 2004. Genomic organization and linkage via a bidirectional promoter of the AP-3 (adaptor protein-3) μ 3A and AK (adenosine kinase) genes: deletion mutants of AK in Chinese hamster cells extend into the AP-3 μ 3A gene. *Biochem. J.* 378: 519-528.
7. Lefrancois, S., et al. 2004. An ear-core interaction regulates the recruitment of the AP-3 complex to membranes. *Dev. Cell* 7: 619-625.
8. Coleman, S.H., et al. 2005. Leucine-specific, functional interactions between human immunodeficiency virus type 1 Nef and adaptor protein complexes. *J. Virol.* 79: 2066-2078.

CHROMOSOMAL LOCATION

Genetic locus: AP3M1 (human) mapping to 10q22.2.

PRODUCT

AP-3 μ siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AP-3 μ shRNA Plasmid (h): sc-60177-SH and AP-3 μ shRNA (h) Lentiviral Particles: sc-60177-V as alternate gene silencing products.

For independent verification of AP-3 μ (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-60177A, sc-60177B and sc-60177C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AP-3 μ siRNA (h) is recommended for the inhibition of AP-3 μ expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AP-3 μ gene expression knockdown using RT-PCR Primer: AP-3 μ (h)-PR: sc-60177-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.