

Agrin (R-20): sc-6040

BACKGROUND

Agrin is a molecule that resides in the basal lamina of muscle cells and directs key events in post synaptic differentiation. Most notably, Agrin is responsible for the clustering of acetylcholine receptors (AChRs) on the cell surface and their localization to the neuromuscular junction. Several Agrin variants have been identified which arise from alternative mRNA splicings. Agrin splice forms having inserts at two sites in the carboxy terminus designated "ψ" and "ζ" display a high affinity for AChRs, while splice forms lacking these inserts associate with AChRs weakly. Muscle α-dystroglycan has been postulated to be the receptor for the clustering activity of agrin; however, this is a point of contention. Tyrosine phosphorylation has been implicated as a required early step in AChR aggregation. Interestingly, a unique receptor tyrosine kinase, designated MuSK, has been discovered that interacts with Agrin and is specifically localized to developing muscle.

REFERENCES

1. Bowen, D.C., et al. 1996. Neural Agrin activates a high-affinity receptor in C2 muscle cells that is unresponsive to muscle Agrin. *J. Neurosci.* 16: 3791-3797.
2. Gautam, M., et al. 1996. Defective neuromuscular synaptogenesis in Agrin-deficient mutant mice. *Cell* 85: 525-535.
3. Slater, C.R. 1996. Agrin signals at the junction. *Nature* 381: 478-479.
4. Gesemann, M., et al. 1996. Alternative splicing of Agrin alters its binding to heparin, dystroglycan, and the putative Agrin receptor. *Neuron* 16: 755-767.
5. O'Toole, J.J., et al. 1996. Alternative splicing of Agrin regulates its binding to heparin α-dystroglycan, and the cell surface. *Proc. Natl. Acad. Sci. USA* 93: 7369-7374.
6. Hopf, C. and Hoch, W. 1996. Agrin binding to α-dystroglycan. Domains of Agrin necessary to induce acetylcholine receptor clustering are overlapping but not identical to the α-dystroglycan-binding region. *J. Biol. Chem.* 271: 5231-5236.
7. Glass, D.J., et al. 1996. Agrin acts via a MuSK receptor complex. *Cell* 85: 513-523.

CHROMOSOMAL LOCATION

Genetic locus: AGRN (human) mapping to 1p36.33; Agrn (mouse) mapping to 4 E2.

SOURCE

Agrin (R-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Agrin of rat origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6040 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Agrin (R-20) is recommended for detection of Agrin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Agrin (R-20) is also recommended for detection of Agrin in additional species, including equine, canine and porcine.

Suitable for use as control antibody for Agrin siRNA (h): sc-29652, Agrin siRNA (m): sc-29653, Agrin shRNA Plasmid (h): sc-29652-SH, Agrin shRNA Plasmid (m): sc-29653-SH, Agrin shRNA (h) Lentiviral Particles: sc-29652-V and Agrin shRNA (m) Lentiviral Particles: sc-29653-V.

Molecular Weight of Agrin: 200 kDa.

Positive Controls: mouse brain extract: sc-2253 or EOC 20 whole cell lysate: sc-364187.

DATA



Agrin (R-20): sc-6040. Western blot analysis of Agrin expression in EOC 20 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Morita, H., et al. 2005. Heparan sulfate of perlecan is involved in glomerular filtration. *J. Am. Soc. Nephrol.* 16: 1703-1710.
2. Tátrai, P., et al. 2006. Agrin, a novel basement membrane component in human and rat liver, accumulates in cirrhosis and hepatocellular carcinoma. *Lab. Invest.* 86: 1149-1160.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Agrin (D-2): sc-374117**, our highly recommended monoclonal alternative to Agrin (R-20). Also, for AC, HRP, FITC, PE, Alexa Fluor[®] 488 and Alexa Fluor[®] 647 conjugates, see **Agrin (D-2): sc-374117**.