

# NEIL2 siRNA (m): sc-61169

## BACKGROUND

NEIL1, NEIL2 and NEIL3, also known as endonuclease VIII-like 1,2 and 3 or DNA-(apurinic or apyrimidinic site) lyase NEIL 1, 2 and 3, are nuclear proteins involved in the repair of DNA damaged by oxidation. The NEIL proteins belong to the FPG family of proteins. They act as DNA glycosylases that can recognize and remove damaged bases, leaving an abasic site. NEIL3, however, lacks the proline residue at the N-terminus which acts as the active site residue found in NEIL1 and NEIL2. NEIL1 is a ubiquitously expressed protein that is up-regulated during S-phase. NEIL2 is expressed primarily in testis, heart, skeletal muscle, placenta, brain, kidney and liver while NEIL3 is detected primarily in thymus and testis.

## REFERENCES

1. Hazra, T.K., et al. 2002. Identification and characterization of a novel human DNA glycosylase for repair of cytosine-derived lesions. *J. Biol. Chem.* 277: 30417-30420.
2. Dou, H., et al. 2003. Repair of oxidized bases in DNA bubble structures by human DNA glycosylases NEIL1 and NEIL2. *J. Biol. Chem.* 278: 49679-49684.
3. Bhakat, K.K., et al. 2004. Acetylation of the human DNA glycosylase NEIL2 and inhibition of its activity. *Nucleic Acids Res.* 32: 3033-3039.
4. Das, A., et al. 2004. Identification of a zinc finger domain in the human NEIL2 (Nei-like-2) protein. *J. Biol. Chem.* 279: 47132-47138.
5. Hailer, M.K., et al. 2005. Recognition of the oxidized lesions spiroiminodihydantoin and guanidinohydantoin in DNA by the mammalian base excision repair glycosylases NEIL1 and NEIL2. *DNA Repair* 4: 41-50.
6. Conlon, K.A., et al. 2005. The murine DNA glycosylase NEIL2 (mNEIL2) and human DNA polymerase  $\beta$  bind microtubules *in situ* and *in vitro*. *DNA Repair* 4: 419-431.

## CHROMOSOMAL LOCATION

Genetic locus: Neil2 (mouse) mapping to 14 D1.

## PRODUCT

NEIL2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NEIL2 shRNA Plasmid (m): sc-61169-SH and NEIL2 shRNA (m) Lentiviral Particles: sc-61169-V as alternate gene silencing products.

For independent verification of NEIL2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-61169A, sc-61169B and sc-61169C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

NEIL2 siRNA (m) is recommended for the inhibition of NEIL2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

NEIL2 (2626C2a): sc-81566 is recommended as a control antibody for monitoring of NEIL2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NEIL2 gene expression knockdown using RT-PCR Primer: NEIL2 (m)-PR: sc-61169-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.