

# RGS10 (N-18): sc-6205

## BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. In mammals, G protein  $\alpha$ ,  $\beta$  and  $\gamma$  poly-peptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their  $\alpha$  subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Four  $G_{\alpha}$  GTPase-activating proteins (GAPs) have been identified and are designated RGS1 (regulator of G protein signaling), RGS4, RGS10 and GAIP ( $G_{\alpha}$ -interacting protein). Each of these proteins has been shown to deactivate specific  $G_{\alpha}$  isoforms by increasing the rate at which they convert GTP to GDP. RGS1, RGS4 and GAIP bind tightly to and exhibit GAP activity towards  $G_{\alpha i}$ ,  $G_{\alpha o}$  and  $G_{\alpha t}$ , but not  $G_{\alpha s}$ . RGS10 increases the GTP hydrolytic activity of several members of the  $G_{\alpha i}$  sub-family, including  $G_{\alpha i-3}$ ,  $G_{\alpha z}$  and  $G_{\alpha o}$ .

## REFERENCES

1. Simon, M.I., et al. 1991. Diversity of G proteins in signal transduction. *Science* 252: 802-808.
2. McLaughlin, S.K., et al. 1992. Gustducin is a taste-cell-specific G protein closely related to the transducins. *Nature* 357: 563-569.

## CHROMOSOMAL LOCATION

Genetic locus: RGS10 (human) mapping to 10q26.11; Rgs10 (mouse) mapping to 7 F3.

## SOURCE

RGS10 (N-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of RGS10 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6205 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

RGS10 (N-18) is recommended for detection of RGS10 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RGS10 siRNA (h): sc-36410, RGS10 siRNA (m): sc-36411, RGS10 shRNA Plasmid (h): sc-36410-SH, RGS10 shRNA Plasmid (m): sc-36411-SH, RGS10 shRNA (h) Lentiviral Particles: sc-36410-V and RGS10 shRNA (m) Lentiviral Particles: sc-36411-V.


Molecular Weight of RGS10: 20 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, BJAB whole cell lysate: sc-2207 or Ramos cell lysate: sc-2216.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



RGS10 (N-18): sc-6205. Western blot analysis of RGS10 expression in BJAB (A) and Jurkat (B) whole cell lysates.



RGS10 (N-18): sc-6205. Immunoperoxidase staining of formalin fixed, paraffin-embedded human tonsil tissue showing cytoplasmic and nuclear staining of cells in germinal center and cells in non-germinal center.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **RGS10 (A-8): sc-46679**, our highly recommended monoclonal alternative to RGS10 (N-18).