

# RGS1 (C-17): sc-6209

## BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. In mammals, G protein  $\alpha$ ,  $\beta$  and  $\gamma$  polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their  $\alpha$  subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Several  $G_{\alpha}$  GTP-ase activating proteins (GAPs) have been identified and are designated RGS1 (regulator of G protein signaling), RGS2, RGS4, RGS7, RGS9, RGS10 and GAIP ( $G_{\alpha}$ -interacting protein). Each of these proteins has been shown to deactivate specific  $G_{\alpha}$  isoforms by increasing the rate at which they convert GTP to GDP. RGS1, RGS4 and GAIP bind tightly to and exhibit GAP activity towards  $G_{\alpha i}$ ,  $G_{\alpha o}$ , and  $G_{\alpha t}$ , but not  $G_{\alpha s}$ .

## REFERENCES

1. Simon, M.I., et al. 1991. Diversity of G proteins in signal transduction. *Science* 252: 802-808.
2. Cali, J.J., et al. 1992. Selective tissue distribution of G protein  $\gamma$  subunits, including a new form of the  $\gamma$  subunits identified by cDNA cloning. *J. Biol. Chem.* 267: 24023-24027.
3. McLaughlin, S.K., et al. 1992. Gustducin is a taste-cell-specific G protein closely related to the transducins. *Nature* 357: 563-569.
4. Kleuss, C., et al. 1992. Different  $\beta$ -subunits determine G-protein interaction with transmembrane receptors. *Nature* 358: 424-426.
5. von Weizsacker, E., et al. 1992. Diversity among the  $\beta$  subunits of heterotrimeric GTP-binding proteins: characterization of a novel  $\beta$ -subunit cDNA. *Biochem. Biophys. Res. Comm.* 183: 350-356.
6. Conklin, B.R., et al. 1993. Structural elements of  $G_{\alpha}$  subunits that interact with  $G_{\beta\gamma}$  receptors, and effectors. *Cell* 73: 631-641.
7. Watson, N., et al. 1996. RGS family members: GTPase-activating proteins for heterotrimeric G-protein  $\alpha$ -subunits. *Nature* 383: 172-175.
8. Heximer, S.P., et al. 1997. RGS2/GOS8 is a selective inhibitor of  $G_{\alpha q}$  function. *Proc. Natl. Acad. Sci. USA* 94: 14389-14393.

## CHROMOSOMAL LOCATION

Genetic locus: RGS1 (human) mapping to 1q31.2; Rgs1 (mouse) mapping to 1 F.

## SOURCE

RGS1 (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of RGS1 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6209 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

RGS1 (C-17) is recommended for detection of RGS1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RGS1 (C-17) is also recommended for detection of RGS1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for RGS1 siRNA (h): sc-36408, RGS1 siRNA (m): sc-36409, RGS1 shRNA Plasmid (h): sc-36408-SH, RGS1 shRNA Plasmid (m): sc-36409-SH, RGS1 shRNA (h) Lentiviral Particles: sc-36408-V and RGS1 shRNA (m) Lentiviral Particles: sc-36409-V.

Molecular Weight of RGS1: 20 kDa.

Positive Controls: BJAB whole cell lysate: sc-2207 or rat intestine extract.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## SELECT PRODUCT CITATIONS

1. Luo, X., et al. 2001. RGS proteins provide biochemical control of agonist-evoked  $[Ca^{2+}]_i$  oscillations. *Mol. Cell* 7: 651-660.
2. Patten, M., et al. 2002. Endotoxin induces desensitization of cardiac endothelin-1 receptor signaling by increased expression of RGS4 and RGS16. *Cardiovasc. Res.* 53: 156-164.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.