

# SCD1/2/3/4 siRNA (m): sc-63288

## BACKGROUND

Stearoyl-CoA desaturase (SCD) is a microsomal enzyme required for the synthesis of oleate and palmitoleate, which are the major monounsaturated fatty acids of membrane phospholipids, triglycerides and cholesterol esters. SCD plays a major role in the triacylglycerol and phospholipid secretion process and in mechanisms of cellular cholesterol homeostasis. It is subject to rapid turnover in the cell and, as such, represents a model for studying selective degradation of short-lived proteins of the ER. SCD is also an important regulator of membrane fluidity. An increase in expression levels of SCD is observed in cells which are induced to differentiate into adipocytes and in certain tumor cell lines. Due to gene duplication events, the number of genes in the SCD family differs between species. Their expression patterns are affected by the level of unsaturated fatty acids in the diet of the animal.

## REFERENCES

1. Ntambi, J.M., et al. 1988. Differentiation-induced gene expression in 3T3-L1 preadipocytes. Characterization of a differentially expressed gene encoding stearoyl-CoA desaturase. *J. Biol. Chem.* 263: 17291-17300.
2. Kaestner, K.H., et al. 1989. Differentiation-induced gene expression in 3T3-L1 preadipocytes. A second differentially expressed gene encoding stearoyl-CoA desaturase. *J. Biol. Chem.* 264: 14755-14761.
3. Li, J., et al. 1994. Partial characterization of a cDNA for human stearoyl-CoA desaturase and changes in its mRNA expression in some normal and malignant tissues. *Int. J. Cancer* 57: 348-352.
4. Miyazaki, M., et al. 2000. The biosynthesis of hepatic cholesterol esters and triglycerides is impaired in mice with a disruption of the gene for stearoyl-coA desaturase 1. *J. Biol. Chem.* 275: 30132-30138.
5. Mziat, H., et al. 2000. The N terminus of microsomal  $\delta$  9 stearoyl-CoA desaturase contains the seunce dterminant for its rapid degradation. *Proc. Natl. Acad. Sci. USA* 97: 8883-8888.
6. Kim, Y.C., et al. 2000. Differential regulation of the stearoyl-CoA desaturase genes by thiazolidinediones in 3T3-L1 adipocytes. *J. Lipid Res.* 41: 1310-1316.

## CHROMOSOMAL LOCATION

Genetic locus: Scd1/Scd2/Scd3/Scd4 (mouse) mapping to 19 C3.

## PRODUCT

SCD1/2/3/4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SCD1/2/3/4 shRNA Plasmid (m): sc-63288-SH and SCD1/2/3/4 shRNA (m) Lentiviral Particles: sc-63288-V as alternate gene silencing products.

For independent verification of SCD1/2/3/4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-63288A, sc-63288B and sc-63288C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

SCD1/2/3/4 siRNA (m) is recommended for the inhibition of SCD1/2/3/4 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

SCD (CD.E10): sc-58420 is recommended as a control antibody for monitoring of SCD1/2/3/4 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SCD1/2/3/4 gene expression knockdown using RT-PCR Primer: SCD1/2/3/4 (m)-PR: sc-63288-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.