

Glucosidase II β (C-20): sc-6451

BACKGROUND

Trimming of glucoses from N-linked core glycans on newly synthesized glycoproteins occurs sequentially through the action of Glucosidases I and II in the endoplasmic reticulum (ER). Glucosidase II is an ER-localized enzyme that contains α and β subunits (Glucosidase II α and Glucosidase II β). The α and β subunits form a defined heterodimeric complex with a molecular weight about 161 kDa. Glucosidase II α is the catalytic core of the enzyme and can function independently of the β subunit. The sequence of Glucosidase II β encodes protein rich in glutamic and aspartic acid with a putative ER retention signal (HDEL) at the C terminus. The phosphorylated form of Glucosidase II β is localized in the plasma membrane and is highly expressed in FGF stimulated fibroblasts and epidermal carcinoma cells. Glucosidase II β was first purified from a human carcinoma cell line as a potential substrate for protein kinase C. Through the HDEL signal at the C-terminus, Glucosidase II β retains the complete complex in the ER.

REFERENCES

1. Shailubhai, K., et al. 1987. Purification and characterization of Glucosidase I involved in N-linked glycoprotein processing in bovine mammary gland. *Biochem. J.* 247: 555-562.
2. Saxena, S., et al. 1987. Purification and characterization of Glucosidase II involved in N-linked glycoprotein processing in bovine mammary gland. *Biochem. J.* 247: 563-570.
3. Trombetta, E.S., et al. 1996. Endoplasmic reticulum Glucosidase II is composed of a catalytic subunit, conserved from yeast to mammals, and a tightly bound noncatalytic HDEL-containing subunit. *J. Biol. Chem.* 271: 27509-27516.
4. Arendt, C.W., et al. 1997. Identification of the CD45-associated 116 kDa and 80 kDa proteins as the α and β subunits of α -Glucosidase II. *J. Biol. Chem.* 272: 13117-13125.
5. Trembl, K., et al. 2000. The α and β subunits are required for expression of catalytic activity in the heterodimeric Glucosidase II complex from human liver. *Glycobiology* 10: 493-502.
6. Trombetta, E.S., et al. 2001. Quaternary and domain structure of glycoprotein processing Glucosidase II. *Biochemistry* 40: 10717-10722.

CHROMOSOMAL LOCATION

Genetic locus: PRKCSH (human) mapping to 19p13.2; Prkcsh (mouse) mapping to 9 A3.

SOURCE

Glucosidase II β (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of Glucosidase II β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6451 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Glucosidase II β (C-20) is recommended for detection of the β subunit of Glucosidase II of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Glucosidase II β siRNA (h): sc-29598, Glucosidase II β siRNA (m): sc-29599, Glucosidase II β shRNA Plasmid (h): sc-29598-SH, Glucosidase II β shRNA Plasmid (m): sc-29599-SH, Glucosidase II β shRNA (h) Lentiviral Particles: sc-29598-V and Glucosidase II β shRNA (m) Lentiviral Particles: sc-29599-V.


Molecular Weight of Glucosidase II β : 80-87 kDa.

Positive Controls: human lung tumor, HeLa whole cell lysate: sc-2200 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Glucosidase II β (C-20): sc-6451. Western blot analysis of Glucosidase II β expression in non-transfected: sc-110760 (A) and human Glucosidase II β transfected: sc-111253 (B) 293 whole cell lysates.



Glucosidase II β (C-20): sc-6451. Western blot analysis of Glucosidase II β expression in non-transfected 293T: sc-117752 (A), mouse Glucosidase II β transfected 293T: sc-120511 (B) and K-562 (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.