



## Rif1 (E-20): sc-65191

### BACKGROUND

Rif1 (Rap1-interacting factor 1) is the 2,472 amino acid homolog of the yeast Rap1 protein and is highly expressed in testis, with increased expression in late G<sub>2</sub>/S phase of the cell cycle. Localized to midzone microtubules during anaphase and to condensed chromosomes during telophase, Rif1 is required for DNA-damage-induced, checkpoint-mediated cell cycle arrest during S phase. The yeast homolog of Rif1 functions by localizing to DNA-damaged foci and binding to uncapped telomeres, thereby inhibiting telomere elongation and slowing cell cycle progression. Human Rif1, unlike its yeast counterpart, does not participate in telomere maintenance or capping, but rather is thought to function at the microtubule midzone in a more global DNA damage response pathway. Rif1 may act by controlling transcription of telomere-related genes or by controlling resolution of twisted chromosomes by topoisomerase II (Topo II). Two isoforms of Rif1 exist due to alternative splicing events.

### REFERENCES

- Li, B. and de Lange, T. 2003. Rap1 affects the length and heterogeneity of human telomeres. *Mol. Biol. Cell.* 14: 5060-5068.
- Silverman, J., Takai, H., Buonomo, S.B., Eisenhaber, F. and de Lange, T. 2004. Human Rif1, ortholog of a yeast telomeric protein, is regulated by Atm and 53BP1 and functions in the S-phase checkpoint. *Genes Dev.* 18: 2108-2119.
- Banerjee, S. and Myung, K. 2004. Increased genome instability and telomere length in the elg1-deficient *Saccharomyces cerevisiae* mutant are regulated by S-phase checkpoints. *Eukaryotic Cell* 3: 1557-1566.
- Xu, L. and Blackburn, E.H. 2004. Human Rif1 protein binds aberrant telomeres and aligns along anaphase midzone microtubules. *J. Cell Biol.* 167: 819-830.
- Hsu, S.Y., Semyonov, J., Park, J.I. and Chang, C.L. 2005. Evolution of the signaling system in relaxin-family peptides. *Ann. N.Y. Acad. Sci.* 1041: 520-529.
- Ji, H., Platts, M.H., Dharamsi, L.M. and Friedman, K.L. 2005. Regulation of telomere length by an N-terminal region of the yeast telomerase reverse transcriptase. *Mol. Cell. Biol.* 25: 9103-9114.
- Miller, K.M., Ferreira, M.G. and Cooper, J.P. 2005. Taz1, Rap1 and Rif1 act both interdependently and independently to maintain telomeres. *EMBO J.* 24: 3128-3135.
- Tsai, H.J., Huang, W.H., Li, T.K., Tsai, Y.L., Wu, K.J., Tseng, S.F. and Teng, S.C. 2006. Involvement of topoisomerase III in telomere-telomere recombination. *J. Biol. Chem.* 281: 13717-13723.

### CHROMOSOMAL LOCATION

Genetic locus: Rif1 (mouse) mapping to 2 C1.1.

### SOURCE

Rif1 (E-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Rif1 of mouse origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-65191 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

Rif1 (E-20) is recommended for detection of Rif1 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Rif1 siRNA (m): sc-62945.

Molecular Weight of Rif1: 274 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.