

UCP2 (N-19): sc-6526

BACKGROUND

The uncoupling protein UCP1 (formerly designated UCP) is an integral membrane protein unique to brown adipose tissue mitochondria. UCP1 forms a dimer that acts as a proton channel, which can uncouple oxidative phosphorylation by dissipating the electrochemical potential across the inner mitochondrial membrane. This process induces heat production in brown adipose tissue and is involved in regulation of body temperature and glucose metabolism. UCP2 is a structurally related protein that also uncouples mitochondrial respiration. It is more widely expressed in human and mouse tissues, including white adipose tissue and muscle, than is UCP. UCP2 is thought to play a role in body weight regulation.

CHROMOSOMAL LOCATION

Genetic locus: UCP2/UCP3 (human) mapping to 11q13.4; Ucp2/Ucp3 (mouse) mapping to 7 E3.

SOURCE

UCP2 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of UCP2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6526 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

UCP2 (N-19) is recommended for detection of UCP2 and, to a lesser extent, UCP3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

UCP2 (N-19) is also recommended for detection of UCP2 and, to a lesser extent, UCP3 in additional species, including equine, canine, bovine and porcine.

Molecular Weight of UCP2 monomer: 35 kDa.

Molecular Weight of UCP2 dimer: 70 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

SELECT PRODUCT CITATIONS

1. Stuart, J.A., et al. 2001. Physiological levels of mammalian uncoupling protein 2 do not uncouple yeast mitochondria. *J. Biol. Chem.* 276: 18633-18639.
2. Bravo, C., et al. 2001. Metabolic changes induced by cold stress in rat liver mitochondria. *J. Bioenerg. Biomembr.* 33: 289-301.
3. Choi, C.S. and Savage, D. 2007. Suppression of diacylglycerol acyltransferase-2 (DGAT2), but not DGAT1, with antisense oligonucleotides reverses diet-induced hepatic steatosis and Insulin resistance. *J. Biol. Chem.* 282: 22678-22688.
4. Teodoro, J.S., et al. 2008. Differential alterations in mitochondrial function induced by a choline-deficient diet: understanding fatty liver disease progression. *Mitochondrion* 8: 367-376.
5. Giardina, T.M., et al. 2008. Uncoupling protein-2 accumulates rapidly in the inner mitochondrial membrane during mitochondrial reactive oxygen stress in macrophages. *Biochim. Biophys. Acta* 1777: 118-129.
6. Selimovic, D., et al. 2008. Taxol-induced mitochondrial stress in melanoma cells is mediated by activation of c-Jun N-terminal kinase (JNK) and p38 pathways via uncoupling protein 2. *Cell. Signal.* 20: 311-322.
7. Amaral, S., et al. 2008. Testicular aging involves mitochondrial dysfunction as well as an increase in UCP2 levels and proton leak. *FEBS Lett.* 582: 4191-4196.
8. Diao, J., et al. 2008. UCP2 is highly expressed in pancreatic α -cells and influences secretion and survival. *Proc. Natl. Acad. Sci. USA* 105: 12057-12062.
9. Liu, Y., et al. 2009. Both ischemic preconditioning and ghrelin administration protect hippocampus from ischemia/reperfusion and upregulate uncoupling protein-2. *BMC Physiol.* 9: 17.
10. Nakaoka, F., et al. 2010. Anti-diabetic effects of globin digest and its active ingredient Leu-Ser-Glu-Leu in ICR mice, streptozotocin-induced diabetic mice and KK-Ay mice. *Life Sci.* 86: 424-434.
11. Jiang, L., et al. 2010. Activation of PPAR α promotes mitochondrial energy metabolism and decreases basal Insulin secretion in palmitate-treated β -cells. *Mol. Cell. Biochem.* 343: 249-256.
12. Calegari, V.C., et al. 2011. Inflammation of the hypothalamus leads to defective pancreatic islet function. *J. Biol. Chem.* 286: 12870-12880.
13. Mailloux, R.J., et al. 2011. Glutathionylation acts as a control switch for uncoupling proteins UCP2 and UCP3. *J. Biol. Chem.* 286: 21865-21875.


 MONOS
 Satisfaction
 Guaranteed

Try **UCP2 (G-6): sc-390189**, our highly recommended monoclonal alternative to UCP2 (N-19). Also, for AC, HRP, FITC, PE, Alexa Fluor[®] 488 and Alexa Fluor[®] 647 conjugates, see **UCP2 (G-6): sc-390189**.