

## FAS (5E2): sc-65274

### BACKGROUND

Cytotoxic T lymphocyte (CTL)-mediated cytotoxicity constitutes an important component of specific effector mechanisms in immunosurveillance against virus-infected or transformed cells. Two mechanisms appear to account for this activity, one of which is the perforin-based process. Independently, a FAS-based mechanism involves the transducing molecule FAS (APO-1) and its ligand (FAS-L). The human FAS (APO-1) protein is a cell surface glycoprotein that belongs to a family of receptors that includes CD40, nerve growth factor receptors and tumor necrosis factor receptors. The FAS antigen is expressed on a broad range of lymphoid cell lines, certain of which undergo apoptosis in response to treatment with antibody to FAS. These findings strongly imply that targeted cell death is potentially mediated by the inter-cellular interactions of FAS with its ligand or effectors, and that FAS may be critically involved in CTL-mediated cytotoxicity.

### REFERENCES

- Henkart, P.A. 1985. Mechanism of lymphocyte-mediated cytotoxicity. *Annu. Rev. Immunol.* 3: 31-58.
- Young, J.D.E., et al. 1988. Perforin-dependent and independent pathways of cytotoxicity mediated by lymphocytes. *Immunol. Rev.* 103: 161-202.
- Podack, E.R., et al. 1991. A central role of perforin in cytolysis? *Annu. Rev. Immunol.* 9: 129-157.
- Yagita, H., et al. 1992. Role of perforin in lymphocyte-mediated cytolysis. *Adv. Immunol.* 51: 215-242.
- Drappa, J., et al. 1993. The FAS protein is expressed at high levels on CD4<sup>+</sup>CD8<sup>+</sup> thymocytes and activated mature lymphocytes in normal mice but not in the lupus-prone strain, MRL lpr/lpr. *Proc. Natl. Acad. Sci. USA* 90: 10340-10344.
- Suda, T., et al. 1993. Molecular cloning and expression of the FAS ligand, a novel member of the tumor necrosis factor family. *Cell* 75: 1169-1178.
- Hanabuchi, S., et al. 1994. FAS and its ligand in a general mechanism of T cell-mediated cytotoxicity. *Proc. Natl. Acad. Sci. USA* 91: 4930-4934.
- Fülöp, P., et al. 2006. Lack of UCP2 reduces FAS-mediated liver injury in ob/ob mice and reveals importance of cell-specific UCP2 expression. *Hepatology* 44: 592-601.
- Maedler, K., et al. 2006. Aging correlates with decreased  $\beta$  cell proliferative capacity and enhanced sensitivity to apoptosis: a potential role for FAS and pancreatic duodenal homeobox-1. *Diabetes* 55: 2455-2462.

### CHROMOSOMAL LOCATION

Genetic locus: FAS (human) mapping to 10q24.1.

### SOURCE

FAS (5E2) is a mouse monoclonal antibody raised against soluble FAS of human origin.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PRODUCT

Each vial contains 100  $\mu$ g IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### APPLICATIONS

FAS (5E2) is recommended for detection of FAS of human origin by immunoprecipitation [1–2  $\mu$ g per 100–500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1  $\mu$ g per 1 x 10<sup>6</sup> cells).


Suitable for use as control antibody for FAS siRNA (h): sc-29311 and FAS siRNA (h2): sc-44260.

Molecular Weight of FAS: 48 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### DATA



FAS (5E2): sc-65274. Indirect FCM analysis of HuT78 cells stained with FAS (5E2), followed by PE-conjugated goat anti-mouse IgG: sc-3738. Black line histogram represents the isotype control, normal mouse IgG<sub>1</sub>: sc-3877.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.