



IL-15 (B-E29): sc-65324

BACKGROUND

Interleukin-15 (IL-15), also designated IL-T, is a cloned cytokine which shares several biological activities but no sequence homology with IL-2. Human, mouse and simian IL-15 cDNA clones have been isolated and characterized. All three species encode a 162 amino acid residue precursor protein containing a 48 amino acid leader that is cleaved to generate the mature form of IL-15. IL-15 stimulates the proliferation of T cells and NK cells, while enhancing B cell expansion and antibody production. Unlike IL-2, IL-15 is not produced by lymphocytes, but appears to be produced by macrophages, epithelial lines, muscle and placenta. IL-15 has also been shown to be a chemoattractant for human blood T lymphocytes and to be able to induce lymphokine-activated killer (LAK) activity in NK cells as well as to be able to induce the generation of cytolytic effector cells. Studies have shown that IL-15 is the only other cytokine that shares the β signaling subunit of the IL-2R. Evidence also suggests that like IL-2, IL-4 and IL-7, IL-15 utilizes the common IL-2R γ subunit.

REFERENCES

- Burton, J.D., Bamford, R.N., Peters, C., Grant, A.J., Kurys, G., Goldman, C.K., Brennan, J., Roessler, E. and Waldmann, T.A. 1994. A lymphokine, provisionally designated interleukin T and produced by a human adult T cell leukemia line, stimulates T cell proliferation and the induction of lymphokine-activated killer cells. *Proc. Natl. Acad. Sci. USA* 91: 4935-4939.
- Grabstein, K.H., Eisenman, J., Shanebeck, K., Rauch, C., Srinivasan, S., Fung, V., Beers, C., Richardson, J., Schoenborn, M.A., Ahdieh, M., Johnson, L., Alderson, M.R., Watson, J.D., Anderson, D.M. and Giri, J.G. 1994. Cloning of a T cell growth factor that interacts with the β chain of the interleukin-2 receptor. *Science* 264: 965-968.
- Giri, J.G., Ahdieh, M., Eisenman, J., Shanebeck, K., Grabstein, K., Kumaki, S., Namen, A., Park, L.S., Cosman, D. and Anderson, D. 1994. Utilization of the β and γ chains of the IL-2 receptor by the novel cytokine IL-15. *EMBO J.* 13: 2822-2830.
- Anderson, D.M., Kumaki, S., Ahdieh, M., Bertles, J., Tometsko, M., Loomis, A., Giri, J., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Valentine, V., Shapiro, D.N., Morris, S.W., Park, L.S. and Cosman, D. 1995. Functional characterization of the human interleukin-15 receptor α chain and close linkage of IL15RA and IL2RA genes. *J. Biol. Chem.* 270: 29862-29869.
- Armitage, R.J., Macduff, B.M., Eisenman, J., Paxton, R. and Grabstein, K.H. 1995. IL-15 has stimulatory activity for the induction of B cell proliferation and differentiation. *J. Immunol.* 154: 483-490.
- Mohamadzadeh, M., Takashima, A., Dougherty, I., Knop, J., Bergstresser, P.R. and Cruz, P.D., Jr. 1995. Ultraviolet B radiation upregulates the expression of IL-15 in human skin. *J. Immunol.* 155: 4492-4496.
- Doherty, T.M., Seder, R.A., and Sher, A. 1996. Induction and regulation of IL-15 expression in murine macrophages. *J. Immunol.* 156: 735-741.

CHROMOSOMAL LOCATION

Genetic locus: IL15 (human) mapping to 4q31; Il15 (mouse) mapping to 8 C2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

IL-15 (B-E29) is a mouse monoclonal antibody raised against recombinant IL-15 of human origin.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

IL-15 (B-E29) is recommended for detection of IL-15 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and flow cytometry (1 μ g per 1×10^6 cells).

Suitable for use as control antibody for IL-15 siRNA (h): sc-39645.

Molecular Weight of IL-15: 14-15 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.