



FAS (2R2): sc-65399

BACKGROUND

Cytotoxic T lymphocyte (CTL)-mediated cytotoxicity constitutes an important component of specific effector mechanisms in immunosurveillance against virus-infected or transformed cells. Two mechanisms appear to account for this activity, one of which is the perforin-based process. Independently, a FAS-based mechanism involves the transducing molecule FAS (also designated APO-1) and its ligand (FAS-L). The human FAS protein is a cell surface glycoprotein that belongs to a family of receptors that includes CD40, nerve growth factor receptors and tumor necrosis factor receptors. The FAS antigen is expressed on a broad range of lymphoid cell lines, certain of which undergo apoptosis in response to treatment with antibody to FAS. These findings strongly imply that targeted cell death is potentially mediated by the intercellular interactions of FAS with its ligand or effectors, and that FAS may be critically involved in CTL-mediated cytotoxicity.

REFERENCES

- Henkart, P.A. 1985. Mechanism of lymphocyte-mediated cytotoxicity. *Ann. Rev. Immunol.* 3: 31-58.
- Young, J.D., Liu, C.C., Persechini, P.M. and Cohn, Z.A. 1988. Perforin-dependent and independent pathways of cytotoxicity mediated by lymphocytes. *Immunol. Rev.* 103: 161-202.
- Podack, E.R., Hengartner, H. and Lichtenheld, M.G. 1991. A central role of perforin in cytolysis? *Ann. Rev. Immunol.* 9: 129-157.
- Yagita, H., Nakata, M., Kawasaki, A., Shinkai, Y. and Okumura, K. 1992. Role of perforin in lymphocyte-mediated cytolysis. *Adv. Immunol.* 51: 215-242.
- Drappa, J., Brot, N. and Elkon, K.B. 1993. The Fas protein is expressed at high levels on CD4⁺CD8⁺ thymocytes and activated mature lymphocytes in normal mice but not in the lupus-prone strain, MRL lpr/lpr. *Proc. Natl. Acad. Sci. USA* 90: 10340-10344.
- Suda, T., Takahashi, T., Golstein, P. and Nagata, S. 1993. Molecular cloning and expression of the Fas ligand, a novel member of the tumor necrosis factor family. *Cell* 75: 1169-1178.
- Hanabuchi, S., Koyanagi, M., Kawasaki, A., Shinohara, N., Matsuzawa, A., Nishimura, Y., Kobayashi, Y., Yonehara, S., Yagita, H. and Okumura, K. 1994. Fas and its ligand in a general mechanism of T cell-mediated cytotoxicity. *Proc. Natl. Acad. Sci. USA* 91: 4930-4934.
- Fulop, P., Dordák, Z., Sheets, A., Sabo, E., Berthiaume, E.P., Resnick, M.B., Wands, J.R., Paragh, G. and Baffy, G. 2006. Lack of UCP2 reduces Fas-mediated liver injury in ob/ob mice and reveals importance of cell-specific UCP2 expression. *Hepatology* 44: 592-601.
- Maedler, K., Schumann, D.M., Schulthess, F., Oberholzer, J., Bosco, D., Berney, T. and Donath, M.Y. 2006. Aging correlates with decreased β -cell proliferative capacity and enhanced sensitivity to apoptosis: a potential role for Fas and pancreatic duodenal homeobox-1. *Diabetes* 55: 2455-2462.

CHROMOSOMAL LOCATION

Genetic locus: FAS (human) mapping to 10q24.1; Fas (mouse) mapping to 19 C1.

SOURCE

FAS (2R2) is a mouse monoclonal antibody raised against FAS of human origin.

PRODUCT

Each vial contains 100 μ g IgG₃ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

FAS (2R2) is recommended for detection of FAS of human origin by flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for FAS siRNA (h): sc-29311 and FAS siRNA (h2): sc-44260.

Molecular Weight of FAS: 48 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.