

# Integrin $\alpha$ M (LM2/1): sc-65429

## BACKGROUND

Integrin  $\alpha$ M (also designated complement component receptor-3  $\alpha$ , CD11b (p170), macrophage antigen a polypeptide, cell surface glycoprotein Mac-1 a subunit, MAC1A, MO1A and ITGAM) is a cell adhesion molecule that acts as a receptor for cell surface ligands such as intracellular adhesion molecules (ICAMs) or soluble ligands. Integrins are heterodimeric proteins that contain an  $\alpha$  chain and  $\beta$  chain. Integrin  $\alpha$ M combines with the Integrin  $\beta$ 2 to form a leukocyte-specific integrin referred to as macrophage receptor 1 (Mac-1), or inactivated-C3b (iC3b) receptor 3 (CR3). Integrin  $\alpha$ M/ $\beta$ 2 is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles.

## REFERENCES

- Nathan, C., et al. 1990. Tumor necrosis factor and CD11/CD18 ( $\beta$ 2) Integrins wact synergistically to lower cAMP in human neutrophils. J. Cell Biol. 111: 2171-2181.
- Li, R., et al. 1995. A peptide derived from the intercellular adhesion molecule-2 regulates the avidity of the leukocyte integrins CD11b/CD18 and CD11c/CD18. J. Cell Biol. 129: 1143-1153.
- Nueda, A., et al. 1995. Hematopoietic cell-type-dependent regulation of leukocyte integrin functional activity: CD11b and CD11c expression inhibits LFA-1-dependent aggregation of differentiated U937 cells. Cell. Immunol. 164: 163-169.
- Walzog, B., et al. 1995. The leukocyte integrin Mac-1 (CD11b/CD18) contributes to binding of human granulocytes to collagen. Exp. Cell Res. 218: 28-38.
- Schlecht, G., et al. 2004. Antigen targeting to CD11b allows efficient presentation of CD4<sup>+</sup> and CD8<sup>+</sup> T cell epitopes and *in vivo* Th1-polarized T cell priming. J. Immunol. 173: 6089-6097.
- Lau, D., et al. 2005. Myeloperoxidase mediates neutrophil activation by association with CD11b/CD18 integrins. Proc. Natl. Acad. Sci. USA 102: 431-436.
- Sandilands, G.P., et al. 2005. Cross-linking of neutrophil CD11b results in rapid cell surface expression of molecules required for antigen presentation and T cell activation. Immunology 114: 354-368.
- Hieronimus, T., et al. 2005. Progressive and controlled development of mouse dendritic cells from Flt3<sup>+</sup>CD11b<sup>+</sup> progenitors *in vitro*. J. Immunol. 174: 2552-2562.
- Carrigan, S.O., et al. 2005. Neutrophil differentiated HL-60 cells model Mac-1 (CD11b/CD18)-independent neutrophil transepithelial migration. Immunology 115: 108-117.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: ITGAM (human) mapping to 16p11.2.

## SOURCE

Integrin  $\alpha$ M (LM2/1) is a mouse monoclonal antibody raised against the I domain of Integrin  $\alpha$ M of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Integrin  $\alpha$ M (LM2/1) is available conjugated to either phycoerythrin (sc-65429 PE) or fluorescein (sc-65429 FITC), 200  $\mu$ g/ml, for WB (RGB), IF, IHC(P) and FCM.

## APPLICATIONS

Integrin  $\alpha$ M (LM2/1) is recommended for detection of Integrin  $\alpha$ M of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1  $\mu$ g per 1 x 10<sup>6</sup> cells).

Suitable for use as control antibody for Integrin  $\alpha$ M siRNA (h): sc-37261, Integrin  $\alpha$ M shRNA Plasmid (h): sc-37261-SH and Integrin  $\alpha$ M shRNA (h) Lentiviral Particles: sc-37261-V.

Molecular Weight of Integrin  $\alpha$ M: 170 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## SELECT PRODUCT CITATIONS

- Day, C.J., et al. 2022. Complement receptor 3 mediates HIV-1 transcytosis across an intact cervical epithelial cell barrier: new insight into HIV transmission in women. mBio. E-published.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **Integrin  $\alpha$ M (2LPM19c): sc-20050** for Integrin  $\alpha$ M antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor<sup>®</sup> 488, 546, 594, 647, 680 and 790.