

Adenovirus (DV4-1A): sc-65652

BACKGROUND

Adenoviruses are non-enveloped icosahedral viruses of the family *Adenoviridae* that contain a double-stranded, linear, non-segmented DNA genome. The Adenovirus life cycle is separated by the DNA replication process into two phases: an early and a late phase. The early phase genes are responsible for expressing mainly nonstructural, regulatory proteins. The late phase of the Adenovirus life cycle is focused on producing sufficient quantities of structural protein to pack all the genetic material produced by DNA replication. Adenovirus infection in humans is usually characterized by illness of the respiratory system, however, depending on the infecting serotype, they may also cause various other symptoms, such as gastroenteritis, conjunctivitis, cystitis and rash. About 47 different types of adenoviruses have been identified, and about half of them are believed to cause human diseases. Infants and children are most commonly affected by adenoviruses.

REFERENCES

- Gallimore, P.H., Sharp, P.A. and Sambrook, J. 1974. Viral DNA in transformed cells II. A study of the sequences of Adenovirus and DNA in nine lines of transformed rat cells using specific fragments of the viral genome. *J. Mol. Biol.* 89: 49-72.
- Jones, N. and Shenk, T. 1979. An Adenovirus type 5 early gene function regulates expression of other early viral genes. *Proc. Natl. Acad. Sci. USA* 76: 3665-3669.
- Berk, A.J., Lee, F., Harrison, T., Williams, J. and Sharp, P.A. 1979. Pre-early Adenovirus-5 gene product regulates synthesis of early viral messenger RNAs. *Cell* 17: 935-944.
- Harlow, E., Franza, B.R. and Schley, C. 1985. Monoclonal antibodies specific for Adenovirus E1A proteins: extensive heterogeneity in the E1A products. *J. Virol.* 3: 533-546.
- Whyte, P., Buchkovich, K.J., Horowitz, J.M., Friend, S.H., Raybuck, M., Weinberg, R.A. and Harlow, E. 1988. Association between an oncogene and an anti-oncogene: the Adenovirus E1A protein binds to the retinoblastoma susceptibility gene product. *Nature* 334: 124-129.
- Giordano, A., Lee, J.H., Scheppeler, J.A., Herrmann, C., Harlow, E., Deuschle, U., Beach, D. and Franza, B.R., Jr. 1991. Cell cycle regulation of Histone H1 kinase activity associated with the adenoviral protein E1A. *Science* 253: 1271-1275.
- Tsai, L.-H., Harlow, E. and Meyerson, M. 1991. Isolation of the human Cdk2 gene that encodes the cyclin A- and Adenovirus E1A-associated p33 kinase. *Nature* 353: 174-177.
- Bell, S.A., Leclere, M., Gardner, I.A. and MacLachlan, N.J. 2006. Equine Adenovirus-1 infection of hospitalised and healthy foals and horses. *Equine Vet. J.* 38: 379-381.
- Rademaker, H.J., Fallaux, F.J., Van den Wollenberg, D.J., De Jong, R.N., Van der Vliet, P.C. and Hoeben, R.C. 2006. Relaxed template specificity in fowl Adenovirus-1 DNA replication initiation. *J. Gen. Virol.* 87: 553-562.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

Adenovirus (DV4-1A) is a mouse monoclonal antibody raised against Adenovirus type 2.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Adenovirus (DV4-1A) is recommended for detection of Adenovirus type 2 by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.