

TAF I p110 (C-18): sc-6566

BACKGROUND

For gene transcription governed by RNA polymerase I, the human transcription factor SL1 (mouse TIF-IB) directs the assembly of initiation complexes at the promoter. Like TFIID, which directs transcription by RNA polymerase II, SL1/TIF-IB contains the TATA-binding protein (TBP) and a set of TBP-associated factors (TAFs). The three TAF I subunits, hTAF I p110, hTAF I p63 and hTAF p48 (or mouse TAF I p95, TAF I p68 and TAF I p48), are all integral components of SL1/TIF-IB. The mutually exclusive binding of either TAF I or TAF II subunits to TBP is believed to direct the formation of promoter- and RNA polymerase-specific complexes.

REFERENCES

1. Learned, R.M., et al. 1985. Purification and characterization of a transcription factor that confers promoter specificity to human RNA polymerase I. *Mol. Cell. Biol.* 5: 1358-1369.
2. Clos, J., et al. 1986. A purified transcription factor (TIF-IB) binds to essential sequences of the mouse rDNA promoter. *Proc. Natl. Acad. Sci. USA* 83: 604-608.

CHROMOSOMAL LOCATION

Genetic locus: TAF1C (human) mapping to 16q24.1.

SOURCE

TAF I p110 (C-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of TAF I p110 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-6566 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-6566 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TAF I p110 (C-18) is recommended for detection of TAF I p110 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TAF I p110 siRNA (h): sc-38490, TAF I p110 shRNA Plasmid (h): sc-38490-SH and TAF I p110 shRNA (h) Lentiviral Particles: sc-38490-V.

TAF I p110 (C-18) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TAF I p110: 110 kDa.

Positive Controls: TAF I p110 (h): 293T Lysate: sc-114680 or PC-3 nuclear extract: sc-2152.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TAF I p110 (C-18): sc-6566. Western blot analysis of TAF I p110 expression in non-transfected: sc-117752 (A) and human TAF I p110 transfected: sc-114680 (B) 293T whole cell lysates.



TAF I p110 (C-18): sc-6566. Western blot analysis of TAF I p110 expression in PC-3 nuclear extract.

SELECT PRODUCT CITATIONS

1. Shen, C.N., et al. 2000. Molecular basis of transdifferentiation of pancreas to liver. *Nat. Cell Biol.* 2: 879-887.
2. Kao, C.F., et al. 2004. Activation of RNA polymerase I transcription by hepatitis C virus core protein. *J. Biomed. Sci.* 11: 72-94.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **TAF I p110 (D-8): sc-376802** or **TAF I p110 (C-10): sc-374551**, our highly recommended monoclonal alternatives to TAF I p110 (C-18).