

AChR β 1 (118): sc-65787

BACKGROUND

Members of the ligand-gated ion channel receptor family are characterized by their fast transmitting response to neurotransmitters. Two important members of this family are the nicotinic acetylcholine and glutamate receptors, both of which are composed of five homologous subunits forming a transmembrane aqueous pore. These transmembrane receptors change conformation in response to their cognate neurotransmitter. Nicotinic acetylcholine receptors (AChRs) are found at the postsynaptic membrane of the neuromuscular junction and bind acetylcholine molecules, allowing ions to move through the pore. Glutamate receptors are found in the postsynaptic membrane of cells in the central nervous system. The activity that is generated at the synapse by the binding of acetylcholine is terminated by acetylcholinesterase, an enzyme that rapidly hydrolyzes acetylcholine. AChR β 1, also known as CHRNB1, CMS1D, CMS2A, SCCMS or CHRNB1, is a 501 amino acid protein that belongs to the ligand-gated ionic channel family. Defects in the gene encoding AChR β 1 may be the cause of congenital myasthenic syndrome slow-channel type (SCCMS), which is characterized by muscle weakness affecting the axial and limb muscles, the ocular muscles and the facial and bulbar musculature.

REFERENCES

- Alkondon, M., et al. 1988. Acetylcholinesterase reactivators modify the functional properties of the nicotinic acetylcholine receptor ion channel. *J. Pharmacol. Exp. Ther.* 245: 543-556.
- Betz, H. 1990. Ligand-gated ion channels in the brain: the amino acid receptor superfamily. *Neuron* 5: 383-392.
- Baenziger, J.E., et al. 1992. Probing conformational changes in the nicotinic acetylcholine receptor by Fourier transform infrared difference spectroscopy. *Biophys. J.* 62: 64-66.
- Daw, N.W., et al. 1993. The role of NMDA receptors in information processing. *Annu. Rev. Neurosci.* 16: 207-222.
- Unwin, N. 1993. Neurotransmitter action: opening of ligand-gated ion channels. *Cell* 72: 31-41.
- Stevens, C.F. 1993. Quantal release of neurotransmitter and long-term potentiation. *Cell* 72: 55-63.
- Sargent, P.B. 1993. The diversity of neuronal nicotinic acetylcholine receptors. *Annu. Rev. Neurosci.* 16: 403-443.
- Ramirez-Latorre, J., et al. 1996. Functional contributions of α 5 subunit to neuronal acetylcholine receptor channels. *Nature* 380: 347-351.

CHROMOSOMAL LOCATION

Genetic locus: CHRNB1 (human) mapping to 17p13.1; Chrn1 (mouse) mapping to 11 B3.

SOURCE

AChR β 1 (118) is a rat monoclonal antibody raised against denatured, purified AChR of *Torpedo* origin.

PRODUCT

Each vial contains 200 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


AChR β 1 (118) is recommended for detection of nicotinic AChR β 1 of mouse, rat, human, bovine and *Torpedo* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for AChR β 1 siRNA (h): sc-29630, AChR β 1 siRNA (m): sc-29631, AChR β 1 shRNA Plasmid (h): sc-29630-SH, AChR β 1 shRNA Plasmid (m): sc-29631-SH, AChR β 1 shRNA (h) Lentiviral Particles: sc-29630-V and AChR β 1 shRNA (m) Lentiviral Particles: sc-29631-V.

Molecular Weight of AChR β 1: 55 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or KNRK whole cell lysate: sc-2214.

DATA



AChR β 1 (118): sc-65787. Western blot analysis of AChR β 1 expression in A-673 (A), SJRH30 (B), Sol8 (C), KNRK (D) and A-10 (E) whole cell lysates.

AChR β 1 (118): sc-65787. Western blot analysis of AChR β 1 expression in HeLa (A), Jurkat (B), K-562 (C), Hep G2 (D) and NIH/3T3 (E) whole cell lysates.

SELECT PRODUCT CITATIONS

- Zhang, D., et al. 2017. Dexamethasone promotes long-term functional recovery of neuromuscular junction in a murine model of tourniquet-induced ischaemia-reperfusion. *Acta Physiol.* 219: 453-464.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.