

PTG (C-20): sc-6581

BACKGROUND

Protein phosphatase 1 (PP1) is a serine-threonine protein phosphatase that plays a central role in mediating the effects of Insulin on glucose and lipid metabolism. PTG (protein targeting to glycogen) was cloned from 3T3-L1 adipocytes as a protein that binds to the PP1 catalytic subunit. The human homolog of PTG, designated PPP1R5, has been shown to bind to PP1 and to modulate its specificity. PTEN/PPP1R5 shows 42% identity to the glycogen binding subunit, G_L , of rat liver PP1. PTG is expressed predominantly in Insulin-sensitive tissues, and it localizes PP1 to glycogen. PTG also has been shown to interact with several enzymes involved in the hormonal regulation of glycogen metabolism, including phosphorylase kinase, phosphorylase A and glycogen synthase. These data indicate a role for PTG in glycogen metabolism, possibly that of a molecular scaffold.

REFERENCES

- Cohen, P. 1989. The structure and regulation of protein phosphatases. *Annu. Rev. Biochem.* 58: 453-508.
- Saltiel, A.R. 1996. Diverse signaling pathways in the cellular actions of Insulin. *Am. J. Physiol.* 270: E375-E385.

CHROMOSOMAL LOCATION

Genetic locus: PPP1R3C (human) mapping to 10q23.32; Ppp1r3c (mouse) mapping to 19 C2.

SOURCE

PTG (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of PTG of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6581 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PTG (C-20) is recommended for detection of PTG of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PTG (C-20) is also recommended for detection of PTG in additional species, including canine, bovine and porcine.


Suitable for use as control antibody for PTG siRNA (h): sc-44047, PTG siRNA (m): sc-152578, PTG shRNA Plasmid (h): sc-44047-SH, PTG shRNA Plasmid (m): sc-152578-SH, PTG shRNA (h) Lentiviral Particles: sc-44047-V and PTG shRNA (m) Lentiviral Particles: sc-152578-V.

Positive Controls: mouse skeletal muscle extract: sc-364250.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: for goat primary antibody (sc-6581): use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), for rabbit primary antibody (sc-6581-R): use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: for goat primary antibody (sc-6581): use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941, for rabbit primary antibody (sc-6581-R): use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PTG (C-20): sc-6581. Western blot analysis of PTG expression in mouse skeletal muscle tissue extract.

SELECT PRODUCT CITATIONS

- Jurczak, M.J., et al. 2007. Transgenic overexpression of protein targeting to glycogen markedly increases adipocytic glycogen storage in mice. *Am. J. Physiol. Endocrinol. Metab.* 292: E952-E963.
- Montori-Grau, M., et al. 2011. Differential pattern of glycogen accumulation after protein phosphatase 1 glycogen-targeting subunit PPP1R6 overexpression, compared to PPP1R3C and PPP1R3A, in skeletal muscle cells. *BMC Biochem.* 12: 57.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.