

Protein C (GMA-067): sc-65958

BACKGROUND

Protein C is a vitamin K-dependent plasma protein that is produced in the liver and made up of two polypeptide chains. It is an important anti-coagulant activated by Thrombin bound to an endothelial surface receptor and it enzymatically cleaves activated forms of Factors V and VIII, thus inhibiting blood coagulation. Protein C is similar to the prothrombin group of blood coagulation factors in its primary structure. Normal protein C concentration in human plasma is approximately 1-3 ng/ml and the proenzyme concentration is approximately 3 µg/ml. Protein C deficiency is associated with inherited thrombophilia, a rare genetic disorder that predisposes affected individuals to venous thrombosis and habitual abortion.

REFERENCES

1. Griffin, J.H., Evatt, B., Zimmerman, T.S., Kleiss, A.J. and Wideman, C. 1982. Deficiency of Protein C in congenital thrombotic disease. *J. Clin. Investig.* 68: 1370-1373.
2. Esmon, C.T. 1989. The roles of Protein C and thrombomodulin in the regulation of blood coagulation. *J. Biol. Chem.* 264: 4743-4746.
3. Dahlbäck, B., Carlsson, M. and Svensson, P.J. 1993. Familial thrombophilia due to a previously unrecognized mechanism characterized by poor anticoagulant response to activated Protein C: prediction of a cofactor to activated Protein C. *Proc. Nat. Acad. Sci. USA* 90: 1004-1008.
4. Bertina, R.M., Koeleman, B.P., Koster, T., Rosendaal, F.R., Dirven, R.J., de Ronde, H., van der Velden, P.A. and Reitsma, P.H. 1994. Mutation in blood coagulation Factor V associated with resistance to activated Protein C. *Nature* 369: 64-67.
5. Dahlbäck, B. 1995. Inherited thrombophilia: resistance to activated Protein C as a pathogenic factor of venous thromboembolism. *Blood* 85: 607-614.
6. Preston, F.E., Rosendaal, F.R., Walker, I.D., Briët, E., Berntorp, E., Conard, J., Fontcuberta, J., Makris, M., Mariani, G., Noteboom, W., Pabinger, I., Legnani, C., Scharrer, I., Schulman, S. and van der Meer, F.J. 1996. Increased fetal loss in women with heritable thrombophilia. *Lancet* 348: 913-916.
7. Abu-Amero, K.K., Owaidah, T.M. and Al-Mahed, M. 2006. Severe type I Protein C deficiency with neonatal purpura fulminans due to a novel homozygous mutation in exon 6 of the Protein C gene. *J. Thromb. Haemost.* 4: 1152-1153.
8. Brueckmann, M., Nahrup, A.S., Lang, S., Bertsch, T., Fukudome, K., Liebe, V., Kaden, J.J., Hoffmann, U., Borggreffe, M. and Huhle, G. 2006. Recombinant human activated Protein C upregulates the release of soluble fractalkine from human endothelial cells. *Br. J. Haematol.* 133: 550-557.
9. Esmon, C.T. 2006. Inflammation and the activated Protein C anticoagulant pathway. *Semin. Thromb. Hemost.* 1: 49-60.

CHROMOSOMAL LOCATION

Genetic locus: PROC (human) mapping to 2q14.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

Protein C (GMA-067) is a mouse monoclonal antibody raised against Protein C of human origin, with epitope mapping to Protein C heavy chain.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


Protein C (GMA-067) is recommended for detection of Protein C of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for Protein C siRNA (h): sc-72054, Protein C shRNA Plasmid (h): sc-72054-SH and Protein C shRNA (h) Lentiviral Particles: sc-72054-V.

Molecular Weight of Protein C: 52 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, ECV304 cell lysate: sc-2269 or Hep G2 cell lysate: sc-2227.

DATA



Protein C (GMA-067): sc-65958. Western blot analysis of Protein C expression in K-562 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.