

Glucose Oxidase (101): sc-66043

BACKGROUND

Glucose Oxidase is a dimeric enzyme that binds to β -D-glucose and aids in its oxidation into D-glucono-1, 5-lactone which then hydrolyzes to gluconic acid. Flavin adenine dinucleotide (FAD) is a cofactor to Glucose Oxidase that acts as the initial electron acceptor and is required for this oxidation to occur. Glucose Oxidase is a natural preservative found in honey, where it reduces atmospheric oxygen into hydrogen peroxide which acts as an antibacterial barrier. Glucose Oxidase is also commonly used in biosensors in which it conveys levels of glucose by keeping track of the number of electrons passed through the enzyme. In this application, Glucose Oxidase is connected to an electrode and the resulting charge is measured.

REFERENCES

1. Subers, M.H. and Schepartz, A.I. 1963. The identification of inhibine, the antibacterial factor in honey, as hydrogen peroxide and its origin in a honey glucose-oxidase system. *Biochim. Biophys. Acta* 73: 57-70.
2. Hartnett, A.M., Ingersoll, C.M., Baker, G.A. and Bright, F.V. 1999. Kinetics and thermodynamics of free flavins and the flavin-based redox active site within Glucose Oxidase dissolved in solution or sequestered within a sol-gel-derived glass. *Anal. Chem.* 71: 1215-1224.
3. Zayats, M., Katz, E. and Willner, I. 2002. Electrical contacting of Glucose Oxidase by surface-reconstitution of the apo-protein on a relay-boronic acid-FAD cofactor monolayer. *J. Am. Chem. Soc.* 124: 2120-2121.
4. Roth, J.P. and Klinman, J.P. 2003. Catalysis of electron transfer during activation of O₂ by the flavoprotein Glucose Oxidase. *Proc. Natl. Acad. Sci. USA* 100: 62-67.
5. Tomotani, E.J., das Neves, L.C. and Vítolo, M. 2005. Oxidation of glucose to gluconic acid by Glucose Oxidase in a membrane bioreactor. *Appl. Biochem. Biotechnol.* 121-124: 149-162
6. Clarke, K.G., Johnstone-Robertson, M., Price, B. and Harrison, S.T. 2006. Location of Glucose Oxidase during production by *Aspergillus niger*. *Appl. Microbiol. Biotechnol.* 70: 72-77.
7. Wu, B.Y., Hou, S.H., Yin, F., Li, J., Zhao, Z.X., Huang, J.D. and Chen, Q. 2006. Amperometric glucose biosensor based on layer-by-layer assembly of multilayer films composed of chitosan, gold nanoparticles and Glucose Oxidase modified Pt electrode. *Biosens. Bioelectron.* 22: 838-844.
8. Morales, M.D., Serra, B., Guzmán-Vázquez de Prada, A., Reviejo, A.J. and Pingarrón, J.M. 2007. An electrochemical method for simultaneous detection and identification of *Escherichia coli*, *Staphylococcus aureus* and *Salmonella choleraesuis* using a glucose oxidase-peroxidase composite biosensor. *Analyst* 132: 572-578.
9. Míguez, D.G., Vanag, V.K. and Epstein, I.R. 2007. Fronts and pulses in an enzymatic reaction catalyzed by Glucose Oxidase. *Proc. Natl. Acad. Sci. USA* 104: 6992-6997.

SOURCE

Glucose Oxidase (101) is a mouse monoclonal antibody raised against Glucose Oxidase of *Aspergillus niger* origin.

PRODUCT

Each vial contains 200 μ g IgM in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Glucose Oxidase (101) is recommended for detection of Glucose Oxidases of *Aspergillus niger* origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Glucose Oxidase: 75 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.