



BoNT/A (24A29): sc-66099

BACKGROUND

Botulism is a rare but serious paralytic illness caused by a nerve toxin, which is produced by the bacterium *Clostridium botulinum*. This neuromuscular disorder occurs through an exquisite series of molecular events, ultimately ending with the arrest of acetylcholine release and hence, flaccid paralysis. Botulinum neurotoxin type A, also known as BoNT/A, Bontoxilysin A and Botox, is one of the most toxic substances known to humans. BoNT/A is a Zn²⁺ endopeptidase which selectively cleaves SNAP 25 (synaptosomal-associated protein), a critical component of the exocytotic machinery. Based on its nucleotide sequence, BoNT/A is a protein that exists as a dimer in aqueous solution. It is also an effective therapeutic drug against involuntary muscle disorders and for pain management. BoNT-A develops its muscle-relaxing effect by the inhibition of acetylcholine (ACh) release.

REFERENCES

1. Ahmed, S.A. and Smith, L.A. 2000. Light chain of botulinum A neurotoxin expressed as an inclusion body from a synthetic gene is catalytically and functionally active. *J. Protein Chem.* 19: 475-487.
2. Byrne, M.P. and Smith, L.A. 2000. Development of vaccines for prevention of botulism. *Biochimie* 82: 955-966.
3. Cai, S. and Singh, B.R. 2001. A correlation between differential structural features and the degree of endopeptidase activity of type A botulinum neurotoxin in aqueous solution. *Biochemistry* 40: 4693-4702.
4. Keller, J.E. and Neale, E.A. 2001. The role of the synaptic protein SNAP 25 in the potency of botulinum neurotoxin type A. *J. Biol. Chem.* 276: 13476-13482.
5. von Lindern, J.J., Niederhagen, B., Appel, T., Berge, S. and Reich, R.H. 2001. Type A botulinum toxin for the treatment of hypertrophy of the masseter and temporal muscles: an alternative treatment. *Plast. Reconstr. Surg.* 107: 327-332.
6. Kelm, S., Gerats, G., Chalkiadaki, A. and Hefter, H. 2001. Reduction of pain and muscle spasms by botulinum toxin A. *Nervenarzt* 72: 302-306.
7. Kobayashi, R., Kohda, T., Kataoka, K., Ihara, H., Kozaki, S., Pascual, D.W., Staats, H.F., Kiyono, H., McGhee, J.R. and Fujihashi, K. 2005. A novel neurotoxicoid vaccine prevents mucosal botulism. *J. Immunol.* 174: 2190-2195.

SOURCE

BoNT/A (24A29) is a mouse monoclonal antibody raised against BoNT/A of *Clostridium botulinum* origin.

PRODUCT

Each vial contains 100 µg IgM in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

BoNT/A (24A29) is recommended for detection of BoNT/A of *Clostridium botulinum* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000).

Molecular Weight of BoNT/A: 162 kDa.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.