

TREX-1 (T-14): sc-66295

BACKGROUND

TREX-1 (3' repair exonuclease-1), also known as trophoblast expressed 1, CRV, AGS1, AGS5, DRN3, HERNS or DNase III, is a member of the exonuclease superfamily and belongs to the TREX family. Members of the TREX family are involved in DNA metabolism and repair. TREX-1 is expressed in spleen, liver, thymus, colon, heart, brain and small intestine. It localizes to the nucleus and exists as a homodimer. TREX-1 is an exonuclease and its activity requires magnesium as a cofactor. TREX-1 has a preference for double stranded DNA and functions in the 3'-5' direction yielding nucleoside 5'-phosphates. TREX-1 also interacts with the SET complex (an endoplasmic reticulum (ER)-associated complex) and participates in granzyme A-mediated apoptosis. Mutations or defects in the gene encoding TREX-1 have been associated with a variety of diseases, including systemic lupus erythematosus, chilblain lupus (CHBL), Aicardi-Goutieres syndrome type 1 (AGS1) and type 5 (AGS5) and autosomal dominant retinal vasculopathy with cerebral leukodystrophy (CRV).

REFERENCES

- Masuda, S., et al. 2005. Recruitment of the human TREX complex to mRNA during splicing. *Genes Dev.* 19: 1512-1517.
- Crow, Y.J., et al. 2006. Mutations in the gene encoding the 3'-5' DNA exonuclease TREX-1 cause Aicardi-Goutieres syndrome at the AGS1 locus. *Nat. Genet.* 38: 917-920.
- Chowdhury, D., et al. 2006. The exonuclease TREX-1 is in the SET complex and acts in concert with nm23-H1 to degrade DNA during granzyme A-mediated cell death. *Mol. Cell* 23: 133-142.
- Wang, J.Y. and Edelman, W. 2006. Mismatch repair proteins as sensors of alkylation DNA damage. *Cancer Cell* 9: 417-418.
- Namiki, Y. and Zou, L. 2006. ATRIP associates with replication protein A-coated ssDNA through multiple interactions. *Proc. Natl. Acad. Sci. USA* 103: 580-585.
- Coscoy, L. and Raulet, D.H. 2007. DNA mismanagement leads to immune system oversight. *Cell* 131: 836-838.
- Yang, Y.G., et al. 2007. TREX-1 exonuclease degrades ssDNA to prevent chronic checkpoint activation and autoimmune disease. *Cell* 131: 873-886.
- Lee-Kirsch, M.A., et al. 2007. Mutations in the gene encoding the 3'-5' DNA exonuclease TREX-1 are associated with systemic lupus erythematosus. *Nat. Genet.* 39: 1065-1067.

CHROMOSOMAL LOCATION

Genetic locus: *Trex1* (mouse) mapping to 9 F2.

SOURCE

TREX-1 (T-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of TREX-1 of mouse origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-66295 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-66295 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

TREX-1 (T-14) is recommended for detection of 3' repair exonuclease 1 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TREX-1 siRNA (m): sc-63158, TREX-1 shRNA Plasmid (m): sc-63158-SH and TREX-1 shRNA (m) Lentiviral Particles: sc-63158-V.

TREX-1 (T-14) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of TREX-1: 33 kDa.

Positive Controls: BC₃H1 cell lysate: sc-2299 or NIH/3T3 whole cell lysate: sc-2210.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **TREX-1 (C-11): sc-133112**, our highly recommended monoclonal alternative to TREX-1 (T-14).