# Na $^+$ /K $^+$ -ATPase $\beta$ 3 (A-18): sc-66337



The Power to Question

#### **BACKGROUND**

The ubiquitously expressed sodium/potassium-ATPase (Na+/K+-ATPase) exists as a oligomeric plasma membrane complex that couples the hydrolysis of one molecule of ATP to the importation of three Na+ ions and two K+ ions against their respective electrochemical gradients. As a member of the P-type family of ion motives, Na+/K+-ATPase plays a critical role in maintaining cellular volume, resting membrane potential and Na+-coupled solute transport. Multiple isoforms of three subunits,  $\alpha$ ,  $\beta$  and  $\gamma$ , comprise the Na+/K+-ATPase oligomer. The  $\alpha$  subunit contains the binding sites for ATP and the cations; the glycosylated  $\beta$  subunit ensures correct folding and membrane insertion of the  $\alpha$  subunits. The small  $\gamma$  subunit co-localizes with the  $\alpha$  subunit in nephron segments, where it increases the affinity of Na+/K+-ATPase for ATP. The  $\beta$  subunit, but not the  $\gamma$  subunit, is essential for normal activity of Na+/K+-ATPase.

## **REFERENCES**

- Hardwicke, P.M., et al. 1981. A proteolipid associated with Na+/K+-ATPase is not essential for ATPase activity. Biochem. Biophys. Res. Commun. 102: 250-257.
- 2. Ackermann, U., et al. 1990. Mutual dependence of Na+/K+-ATPase  $\alpha$  and  $\beta$ -subunits for correct post-translational processing and intracellular transport. FEBS Letts. 269: 105-108.
- 3. McDonough, A.A., et al. 1990. The sodium pump needs its  $\beta$  subunit. FASEB J. 4: 1598-1605.
- Pedemonte, C.H., et al. 1990. Chemical modification as an approach to elucidation of sodium pump structure-function relations. Am. J. Physiol. 258: C1-23.
- Mercer, R.W., et al. 1993. Molecular cloning and immunological chracterization of the γ-polypeptide, a small protein associated with Na+/K+-ATPase.
  J. Cell Biol. 121: 579-586.
- DeTomaso, A.W., et al. 1993. Expression, targeting, and assembly of functional Na+/K+-ATPase polypeptides in baculovirus-infected insect cells. J. Biol. Chem. 268: 1470-1478.
- Scheiner-Bobis, G., et al. 1994. Subunit requirements for expression of functional sodium pumps in yeast cells. Biochim. Biophys. Acta 1193: 226-234.
- 8. Lingrel, J.B., et al. 1994. Na+,K+-ATPase. J. Biol. Chem. 269: 19659-19662.
- 9. Blanco, G., et al. 1994. The  $\alpha$ -subunit of the Na+/K+-ATPase has catalytic activity independent of the  $\beta$ -subunit. J. Biol. Chem. 269: 23420-23425.

## CHROMOSOMAL LOCATION

Genetic locus: (mouse) mapping to .

## SOURCE

Na+/K+-ATPase  $\beta 3$  (A-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Na+/K+-ATPase  $\beta 3$  of rat origin.

#### **PRODUCT**

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-66337 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **APPLICATIONS**

Na+/K+-ATPase  $\beta$ 3 (A-18) is recommended for detection of Na+/K+-ATPase  $\beta$ 3 of rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Na+/K+-ATPase β3: 40-60 kDa.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

# **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com