

Na⁺/K⁺-ATPase β3 (I-14): sc-66340

BACKGROUND

The ubiquitously expressed sodium/potassium-ATPase (Na⁺/K⁺-ATPase) exists as a oligomeric plasma membrane complex that couples the hydrolysis of one molecule of ATP to the importation of three Na⁺ ions and two K⁺ ions against their respective electrochemical gradients. As a member of the P-type family of ion motives, Na⁺/K⁺-ATPase plays a critical role in maintaining cellular volume, resting membrane potential and Na⁺-coupled solute transport. Multiple isoforms of three subunits, α, β and γ, comprise the Na⁺/K⁺-ATPase oligomer. The α subunit contains the binding sites for ATP and the cations; the glycosylated β subunit ensures correct folding and membrane insertion of the α subunits. The small γ subunit co-localizes with the α subunit in nephron segments, where it increases the affinity of Na⁺/K⁺-ATPase for ATP. The β subunit, but not the γ subunit, is essential for normal activity of Na⁺/K⁺-ATPase.

REFERENCES

1. Hardwicke, P.M., et al. 1981. A proteolipid associated with Na⁺/K⁺-ATPase is not essential for ATPase activity. *Biochem. Biophys. Res. Commun.* 102: 250-257.
2. Ackermann, U., et al. 1990. Mutual dependence of Na⁺/K⁺-ATPase α- and β-subunits for correct post-translational processing and intracellular transport. *FEBS Lett.* 269: 105-108.
3. McDonough, A.A., et al. 1990. The sodium pump needs its β subunit. *FASEB J.* 4: 1598-1605.
4. Pedemonte, C.H., et al. 1990. Chemical modification as an approach to elucidation of sodium pump structure-function relations. *Am. J. Physiol.* 258: C1-C23.
5. Mercer, R.W., et al. 1993. Molecular cloning and immunological characterization of the γ-polypeptide, a small protein associated with Na⁺/K⁺-ATPase. *J. Cell Biol.* 121: 579-586.
6. DeTomaso, A.W., et al. 1993. Expression, targeting, and assembly of functional Na⁺/K⁺-ATPase polypeptides in baculovirus-infected insect cells. *J. Biol. Chem.* 268: 1470-1478.
7. Scheiner-Bobis, G., et al. 1994. Subunit requirements for expression of functional sodium pumps in yeast cells. *Biochim. Biophys. Acta* 1193: 226-234.
8. Lingrel, J.B., et al. 1994. Na⁺/K⁺-ATPase. *J. Biol. Chem.* 269: 19659-19662.
9. Blanco, G., et al. 1994. The α-subunit of the Na⁺/K⁺-ATPase has catalytic activity independent of the β-subunit. *J. Biol. Chem.* 269: 23420-23425.

CHROMOSOMAL LOCATION

Genetic locus: *Atp1b3* (mouse) mapping to 9 E3.3.

SOURCE

Na⁺/K⁺-ATPase β3 (I-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Na⁺/K⁺-ATPase β3 of mouse origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-66340 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Na⁺/K⁺-ATPase β3 (I-14) is recommended for detection of Na⁺/K⁺-ATPase β3 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Na⁺/K⁺-ATPase β3 siRNA (m): sc-62003, Na⁺/K⁺-ATPase β3 shRNA Plasmid (m): sc-62003-SH and Na⁺/K⁺-ATPase β3 shRNA (m) Lentiviral Particles: sc-62003-V.

Molecular Weight of Na⁺/K⁺-ATPase β3: 40-60 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.