



OATP-B (T-20): sc-66562

BACKGROUND

The organic anion transporter family of proteins mediate hepatic uptake of cardiac glycosides. OATP-A and OATP-C are both pravastatin transporters, suggesting that they are responsible for the hepatic uptake of the liver-specific hydroxymethylglutaryl-CoA reductase inhibitor in mouse, rat and human. OATP-A is expressed in liver and kidney and helps mediate sodium-independent uptake of the anionic steroid conjugates dehydroepiandrosterone sulfate, estradiol-17-glucuronide and prostaglandin. OATP-C is exclusively expressed in liver and is localized to the basolateral hepatocyte membrane. OATP-B, also known as Slco2b1 or Slc21a9, mediates the Na^+ independent transport of organic anions such as taurocholate, leukotriene C₄, thromboxane B₂ and iloprost during the absorption of bile acids in the liver.

REFERENCES

1. Nagase, T., et al. 1999. Prediction of the coding sequences of unidentified human genes. XII. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 5: 355-364.
2. St-Pierre, M.V., et al. 2002. Characterization of an organic anion transporting polypeptide (OATP-B) in human placenta. J. Clin. Endocrinol. Metab. 87: 1856-1863.
3. Pizzagalli, F., et al. 2002. Identification of a novel human organic anion transporting polypeptide as a high affinity thyroxine transporter. Mol. Endocrinol. 16: 2283-2296.
4. Cai, S.Y., et al. 2002. An evolutionarily ancient Oatp: insights into conserved functional domains of these proteins. Am. J. Physiol. Gastrointest. Liver Physiol. 282: G702-710.
5. Takagi, M., et al. 2004. Enhancement of the inhibitory activity of Oatp antisense oligonucleotides by incorporation of 2'-O,4'-C-ethylene-bridged nucleic acids (ENA) without a loss of subtype selectivity. Biochemistry 43: 4501-4510.
6. Niemi, M., et al. 2004. High plasma pravastatin concentrations are associated with single nucleotide polymorphisms and haplotypes of organic anion transporting polypeptide-C (OATP-C, SLCO1B1). Pharmacogenetics 14: 429-440.
7. Satoh, H., et al. 2005. Citrus juices inhibit the function of human organic anion transporting polypeptide OATP-B. Drug Metab. Dispos. 33: 518-523.
8. Kopplow, K., et al. 2005. Human hepatobiliary transport of organic anions analyzed by quadruple-transfected cells. Mol. Pharmacol. 68: 1031-1038.
9. Fuchikami, H., et al. 2006. Effects of herbal extracts on the function of human organic anion transporting polypeptide OATP-B. Drug Metab. Dispos. 34: 577-582.

CHROMOSOMAL LOCATION

Genetic locus: Slco2b1 (mouse) mapping to 7 E1.

SOURCE

OATP-B (T-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of OATP-B of mouse origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-66562 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

OATP-B (T-20) is recommended for detection of OATP-B of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for OATP-B siRNA (m): sc-62712, OATP-B shRNA Plasmid (m): sc-62712-SH and OATP-B shRNA (m) Lentiviral Particles: sc-62712-V.

Molecular Weight of OATP-B: 85 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.