

PON2 (C-13): sc-66578

BACKGROUND

Paroxon is an organophosphorus anticholinesterase compound, used topically in the treatment of glaucoma. It is produced *in vivo* in mammals by microsomal oxidation of the insecticide parathion. Parathion is inert until transformed to paroxon. Paroxonase (paraoxonase or PON) is an arylesterase that is capable of hydrolyzing paroxon to produce p-nitrophenol. PONs are nonspecific and their classification is based not only on substrate specificity but also on tissue distribution, inhibition properties and physicochemical characteristics such as electrophoretic mobility and molecular weight. In contrast to PON1, which is expressed mainly in the liver, PON2 is expressed in a variety of mouse tissues, including the pancreas. PON3 is associated with the high density lipoprotein fraction of serum. The genes which encode PON1-3 are physically linked and map to human chromosome 7q21.3.

REFERENCES

1. Coates, P.M., et al. 1975. A preliminary genetic interpretation of the esterase isozymes of human tissues. *Ann. Hum. Genet.* 39: 1-20.
2. Humbert, R., et al. 1993. The molecular basis of the human serum paraoxonase activity polymorphism. *Nat. Genet.* 3: 73-76.
3. Primo-Parmo, S.L., et al. 1996. The human serum paraoxonase/arylesterase gene (PON1) is one member of a multigene family. *Genomics* 33: 498-507.
4. Mochizuki, H., et al. 1998. Human PON2 gene at 7q21.3: cloning, multiple mRNA forms, and missense polymorphisms in the coding sequence. *Gene* 213: 149-157.
5. Draganov, D.I., et al. 2000. Rabbit serum paraoxonase 3 (PON3) is a high density lipoprotein-associated lactonase and protects low density lipoprotein against oxidation. *J. Biol. Chem.* 275: 33435-33442.
6. LocusLink Report (LocusID: 168820).
<http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: PON2 (human) mapping to 7q21.3.

SOURCE

PON2 (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of PON2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-66578 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

PON2 (C-13) is recommended for detection of PON2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PON2 (C-13) is also recommended for detection of PON2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PON2 siRNA (h): sc-62838, PON2 shRNA Plasmid (h): sc-62838-SH and PON2 shRNA (h) Lentiviral Particles: sc-62838-V.

Molecular Weight of PON2: 40 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **PON2 (C-5): sc-374158** or **PON2 (D-12): sc-373981**, our highly recommended monoclonal alternatives to PON2 (C-13).