

Riboflavin kinase (N-14): sc-66627

BACKGROUND

Riboflavin kinase, also known as RFK or RIFK, is a cytoplasmic protein that catalyzes the first step in flavocoenzyme biosynthesis, namely the ATP-dependent phosphorylation of riboflavin to form flavin-mononucleotide (FMN). Expressed in the brain, placenta and bladder, Riboflavin kinase is a 162 amino acid protein for which zinc and magnesium are cofactors. Riboflavin kinase has three distinct conformational states that are referred to as the binary MgADP complex, the ternary product complex and the apo form, all of which contribute to the unique substrate binding and catalytic activity of the enzyme. Human Riboflavin kinase shares 44% homology with its yeast counterpart, suggesting that the three flexible regions surrounding the active site (termed Flap I, Flap II and Helix B) are similar in both species.

REFERENCES

- Barile, M., Brizio, C., Valenti, D., De Virgilio, C. and Passarella, S. 2000. The riboflavin/FAD cycle in rat liver mitochondria. *Eur. J. Biochem.* 267: 4888-4900.
- Karthikeyan, S., Zhou, Q., Mseeh, F., Grishin, N.V., Osterman, A.L. and Zhang, H. 2003. Crystal structure of human riboflavin kinase reveals a beta barrel fold and a novel active site arch. *Structure* 11: 265-273.
- Karthikeyan, S., Zhou, Q., Osterman, A.L. and Zhang, H. 2003. Ligand binding-induced conformational changes in riboflavin kinase: structural basis for the ordered mechanism. *Biochemistry* 42: 12532-12538.
- Solovieva, I.M., Kreneva, R.A., Errais Lopes, L. and Perumov, D.A. 2005. The riboflavin kinase encoding gene *ribR* of *Bacillus subtilis* is a part of a 10 kb operon, which is negatively regulated by the *yzc* gene product. *FEMS Microbiol. Lett.* 243: 51-58.
- Sandoval, F.J. and Roje, S. 2005. An FMN hydrolase is fused to a riboflavin kinase homolog in plants. *J. Biol. Chem.* 280: 38337-38345.
- Bertollo, C.M., Oliveira, A.C., Rocha, L.T., Costa, K.A. and Coelho, M.M. 2006. Characterization of the antinociceptive and anti-inflammatory activities of riboflavin in different experimental models. *Eur. J. Pharmacol.* 547: 184-191.
- Ammelburg, M., Hartmann, M.D., Djuranovic, S., Alva, V., Koretke, K.K., Martin, J., Sauer, G., Truffault, V., Zeth, K., Lupas, A.N. and Coles, M. 2007. A CTP-dependent archaeal riboflavin kinase forms a bridge in the evolution of cradle-loop barrels. *Structure* 15: 1577-1590.

CHROMOSOMAL LOCATION

Genetic locus: RFK (human) mapping to 9q21.13.

SOURCE

Riboflavin kinase (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Riboflavin kinase of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-66627 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Riboflavin kinase (N-14) is recommended for detection of Riboflavin kinase of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Riboflavin kinase siRNA (h): sc-62940, Riboflavin kinase shRNA Plasmid (h): sc-62940-SH and Riboflavin kinase shRNA (h) Lentiviral Particles: sc-62940-V.


Molecular Weight of Riboflavin kinase: 18 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Riboflavin kinase (N-14): sc-66627. Western blot analysis of Riboflavin kinase expression in HeLa (A) and Jurkat (B) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.