

# COP1 (H-300): sc-66837

## BACKGROUND

COP1 (constitutive photomorphogenesis protein 1), also designated RFWD2 (ring finger and WD repeat domain 2) or RNF200 (ring finger protein 200), is an E3 ubiquitin ligase protein that mediates ubiquitination and degradation of target proteins such as c-Jun and p53. It is a component of the DCX DET1-COP1 ubiquitin ligase complex which consists of RBX1, DET1, DDB1, CUL4A and COP1. Localizing to the cytoplasm and to the nucleus, COP1 is primarily expressed in testis, placenta, heart and skeletal muscle. COP1 is a potent inhibitor of p53-dependent transcription and apoptosis but, when phosphorylated by Atm (ataxia telangiectasia mutated) in response to DNA damage, the COP1-p53 complex is disrupted and p53 is allowed to exert its pro-apoptotic properties. In ovarian and breast cancers, COP1 is overexpressed, suggesting a role for COP1 in tumorigenesis.

## CHROMOSOMAL LOCATION

Genetic locus: RFWD2 (human) mapping to 1q25.1; Rfwd2 (mouse) mapping to 1 H1.

## SOURCE

COP1 (H-300) is a rabbit polyclonal antibody raised against amino acids 432-731 mapping at the C-terminus of COP1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-66837 X, 200 µg/0.1 ml.

## APPLICATIONS

COP1 (H-300) is recommended for detection of COP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000). COP1 (H-300) is also recommended for detection of COP1 in additional species, including equine, canine, bovine and avian.

COP1 (H-300) is also recommended for detection of COP1 in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for COP1 siRNA (h): sc-45541, COP1 siRNA (m): sc-45542, COP1 shRNA Plasmid (h): sc-45541-SH, COP1 shRNA Plasmid (m): sc-45542-SH, COP1 shRNA (h) Lentiviral Particles: sc-45541-V and COP1 shRNA (m) Lentiviral Particles: sc-45542-V.

COP1 (H-300) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight (predicted) of COP1: 80 kDa.


Molecular Weight (observed) of COP1: 115 kDa.

Positive Controls: JAR cell lysate: sc-2276 or T24 cell lysate: sc-2292.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

## DATA



COP1 (H-300): sc-66837. Western blot analysis of COP1 expression in JAR whole cell lysate.



COP1 (H-300): sc-66837. Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing cytoplasmic, membrane and faint nuclear staining of decidual cells.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **COP1 (B-12): sc-166799**, our highly recommended monoclonal alternative to COP1 (H-300).