

# Coronin 1C (H-85): sc-66840

## BACKGROUND

Coronins are a family of WD repeat-containing, actin-binding proteins that localize to submembrane areas and regulate cell motility and cytoskeletal rearrangement. Coronin 1A (CORO1A, CLIPINA, CLABP, TACO, p57) can form coiled coil-mediated homotrimeric complexes that influence early phagosome formation. PKC-dependent phosphorylation of Coronin 1B (CORO1B) at Serine 2 regulates leading edge dynamics and cell motility in fibroblasts through interactions with Arp2/3 complex. Coronin 1C (CORO1C, Coronin 3, HCRNN4) is abundant in differentiating Neuro-2a cells, PC-12 cells and primary oligodendrocytes, where it is thought to influence neuron morphogenesis and migration. Coronin 2A (CORO2A, CLIPINB, IR10, WDR2) is a component of the approximately 1.5-2 megadalton N-CoR (nuclear receptor corepressor) complex of 10-12 proteins, which recruits HDACs to generate repressive chromatin. Coronin 7 (CORO7, CRN7) localizes to the Golgi membrane and influences the organization of intracellular membrane compartments and vesicular trafficking. Coronin 2B (CORO2B, CLIPINC) and Coronin 6 (CORO6) are similar to other members of this family, since they possess a conserved basic N-terminal motif and 3-10 WD repeats clustered in one to two core domains.

## CHROMOSOMAL LOCATION

Genetic locus: CORO1C (human) mapping to 12q24.11; Coro1c (mouse) mapping to 5 F.

## SOURCE

Coronin 1C (H-85) is a rabbit polyclonal antibody raised against amino acids 390-474 mapping at the C-terminus of Coronin 1C of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Coronin 1C (H-85) is recommended for detection of Coronin 1C of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Coronin 1C (H-85) is also recommended for detection of Coronin 1C in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Coronin 1C siRNA (h): sc-44693, Coronin 1C siRNA (m): sc-44694, Coronin 1C shRNA Plasmid (h): sc-44693-SH, Coronin 1C shRNA Plasmid (m): sc-44694-SH, Coronin 1C shRNA (h) Lentiviral Particles: sc-44693-V and Coronin 1C shRNA (m) Lentiviral Particles: sc-44694-V.


Molecular Weight of Coronin 1C: 57 kDa.

Positive Controls: HeLa nuclear extract: sc-2120.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Coronin 1C (H-85): sc-66840. Western blot analysis of Coronin 1C expression in HeLa nuclear extract.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **Coronin 1C (D-9): sc-376919** or **Coronin 1C (G-R2): sc-130448**, our highly recommended monoclonal alternatives to Coronin 1C (H-85).