

Coronin 2A (H-110): sc-66841

BACKGROUND

Coronins are a family of WD repeat-containing, actin-binding proteins that localize to submembrane areas and regulate cell motility and cytoskeletal rearrangement. Coronin 1A (CORO1A, CLIPINA, CLABP, TACO, p57) can form coiled coil-mediated homotrimeric complexes that influence early phagosome formation. PKC-dependent phosphorylation of Coronin 1B (CORO1B) at Serine 2 regulates leading edge dynamics and cell motility in fibroblasts through interactions with Arp2/3 complex. Coronin 1C (CORO1C, Coronin 3, HCRNN4) is abundant in differentiating Neuro-2a cells, PC-12 cells and primary oligodendrocytes, where it is thought to influence neuron morphogenesis and migration. Coronin 2A (CORO2A, CLIPINB, IR10, WDR2) is a component of the approximately 1.5-2 megadalton N-CoR (nuclear receptor corepressor) complex of 10-12 proteins, which recruits HDACs to generate repressive chromatin. Coronin 7 (CORO7, CRN7) localizes to the Golgi membrane and influences the organization of intracellular membrane compartments and vesicular trafficking. Coronin 2B (CORO2B, CLIPINC) and Coronin 6 (CORO6) are similar to other members of this family, since they possess a conserved basic N-terminal motif and three-ten WD repeats clustered in one to two core domains.

REFERENCES

- Mishima, M., et al. 1999. Coronin localizes to leading edges and is involved in cell spreading and lamellipodium extension in vertebrate cells. *J. Cell Sci.* 112: 2833-2842.
- Spoerl, Z., et al. 2002. Oligomerization, F-Actin interaction, and membrane association of the ubiquitous mammalian Coronin 3 are mediated by its carboxyl-terminus. *J. Biol. Chem.* 277: 48858-48867.
- Yoon, H.G., et al. 2003. Purification and functional characterization of the human N-CoR complex: the roles of HDAC3, TBL1 and TBLR1. *EMBO J.* 22: 1336-1346.
- Rybakin, V., et al. 2004. Coronin 7, the mammalian POD-1 homologue, localizes to the Golgi apparatus. *FEBS Lett.* 573: 161-167.
- Gatfield, J., et al. 2005. Association of the leukocyte plasma membrane with the Actin cytoskeleton through coiled coil-mediated trimeric Coronin 1 molecules. *Mol. Biol. Cell* 16: 2786-2798.

CHROMOSOMAL LOCATION

Genetic locus: CORO2A (human) mapping to 9q22.33.

SOURCE

Coronin 2A (H-110) is a rabbit polyclonal antibody raised against amino acids 371-480 mapping near the C-terminus of Coronin 2A of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Coronin 2A (H-110) is recommended for detection of Coronin 2A of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Coronin 2A siRNA (h): sc-44685, Coronin 2A shRNA Plasmid (h): sc-44685-SH and Coronin 2A shRNA (h) Lentiviral Particles: sc-44685-V.


Molecular Weight of Coronin 2A: 57 kDa.

Positive Controls: Coronin 2A (h2): 293T Lysate: sc-172930.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Coronin 2A (H-110): sc-66841. Western blot analysis of Coronin 2A expression in non-transfected: sc-117752 (A) and human Coronin 2A transfected: sc-172930 (B) 293T whole cell lysates.



Coronin 2A (H-110): sc-66841. Western blot analysis of Coronin 2A expression in non-transfected: sc-117752 (A) and human Coronin 2A transfected: sc-172290 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE** Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


 MONOS
Satisfaction
Guaranteed

Try **Coronin 2A (C-7): sc-376197** or **Coronin 2A (H-2): sc-376194**, our highly recommended monoclonal alternatives to Coronin 2A (H-110).