

Cyclophilin 40 (H-185): sc-66848

BACKGROUND

Cyclophilins are conserved, ubiquitous and abundant cytosolic peptidyl-prolyl *cis-trans* isomerases that accelerate the isomerization of XaaPro peptide bonds and the refolding of proteins. Human Cyclophilin A (CyPA), an intracellular protein of 165 amino acids, is the target of Cyclosporin A (CsA) and is encoded by a single unique gene conserved from yeast to humans. Cyclophilin B (CyPB) is secreted in biological fluids such as blood or milk and binds to a specific receptor present on the human lymphoblastic cell line Jurkat and on human peripheral blood lymphocytes. Cyclophilin 40 (CyP40) is a widely expressed cytoplasmic protein that catalyzes the *cis-trans* isomerization of proline imidic peptide bonds in oligopeptides. It is a widely expressed cytoplasmic protein.

REFERENCES

1. Kieffer, L.J., et al. 1993. Cyclophilin 40, a protein with homology to the P59 component of the steroid receptor complex. Cloning of the cDNA and further characterization. *J. Biol. Chem.* 268: 12303-12310.
2. Yokoi, H., et al. 1996. The structure and complete nucleotide sequence of the human Cyclophilin 40 (PPID) gene. *Genomics* 35: 448-455.

CHROMOSOMAL LOCATION

Genetic locus: PPID (human) mapping to 4q32.1; Ppid (mouse) mapping to 3 E3.

SOURCE

Cyclophilin 40 (H-185) is a rabbit polyclonal antibody raised against amino acids 186-370 mapping at the C-terminus of Cyclophilin 40 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Cyclophilin 40 (H-185) is recommended for detection of Cyclophilin 40 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Cyclophilin 40 (H-185) is also recommended for detection of Cyclophilin 40 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Cyclophilin 40 siRNA (h): sc-44892, Cyclophilin 40 siRNA (m): sc-44893, Cyclophilin 40 shRNA Plasmid (h): sc-44892-SH, Cyclophilin 40 shRNA Plasmid (m): sc-44893-SH, Cyclophilin 40 shRNA (h) Lentiviral Particles: sc-44892-V and Cyclophilin 40 shRNA (m) Lentiviral Particles: sc-44893-V.


Molecular Weight of Cyclophilin 40: 41 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210 or JAR cell lysate: sc-2276.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



Cyclophilin 40 (H-185): sc-66848. Western blot analysis of Cyclophilin 40 expression in JAR (A) and NIH/3T3 (B) whole cell lysates.



Cyclophilin 40 (H-185): sc-66848. Immunofluorescence staining of formalin-fixed HepG2 cells showing nuclear and cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human esophagus tissue showing cytoplasmic and nuclear staining of squamous epithelial cells (B).

SELECT PRODUCT CITATIONS

1. Shimamoto, S., et al. 2010. S100 proteins regulate the interaction of Hsp90 with Cyclophilin 40 and FKBP52 through their tetratricopeptide repeats. *FEBS Lett.* 584: 1119-1125.
2. Janevski, M., et al. 2012. Fructose containing sugars modulate mRNA of lipogenic genes ACC and FAS and protein levels of transcription factors ChREBP and SREBP1c with no effect on body weight or liver fat. *Food Funct.* 3: 141-149.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **Cyclophilin 40 (C-11): sc-137216** or **Cyclophilin 40 (F-5): sc-137157**, our highly recommended monoclonal alternatives to Cyclophilin 40 (H-185).