

# ERI-1 (H-135): sc-66881

## BACKGROUND

Helicase with RNase motif, more commonly designated dicer, cleaves double-stranded RNA (dsRNA) in the RNA interference and small temporal RNA (stRNA) pathways, producing active small RNA components (siRNAs) which target the destruction of RNA and repress gene expression. Human dicer cleaves dsRNA independent of ATP. The 3'-5' exonuclease ERI-1, also known as protein 3'hExo, degrades Histone mRNA after replication and may be involved in the regulation of RNA interference. ERI-1 has a high affinity for the stem-loop structure of replication-dependent Histone pre-mRNAs. It requires the 5'-ACCCA-3' sequence present in stem-loop structure. ERI-1 and a stem-loop binding protein (SLBP) target opposite faces of a unique highly conserved stem-loop RNA scaffold towards the 3' end of Histone mRNA.

## REFERENCES

1. Kennedy, S., et al. 2004. A conserved siRNA-degrading RNase negatively regulates RNA interference in *C. elegans*. *Nature* 427: 645-649.
2. Timmons, L. 2004. Endogenous inhibitors of RNA interference in *Caenorhabditis elegans*. *Bioessays* 26: 715-718.

## CHROMOSOMAL LOCATION

Genetic locus: ERI1 (human) mapping to 8p23.1; Eri1 (mouse) mapping to 8 A4.

## SOURCE

ERI-1 (H-135) is a rabbit polyclonal antibody raised against amino acids 161-295 mapping within an internal region of ERI-1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

ERI-1 (H-135) is recommended for detection of ERI-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ERI-1 (H-135) is also recommended for detection of ERI-1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ERI-1 siRNA (h): sc-45559, ERI-1 siRNA (m): sc-45560, ERI-1 shRNA Plasmid (h): sc-45559-SH, ERI-1 shRNA Plasmid (m): sc-45560-SH, ERI-1 shRNA (h) Lentiviral Particles: sc-45559-V and ERI-1 shRNA (m) Lentiviral Particles: sc-45560-V.

Molecular Weight (predicted) of ERI-1: 34 kDa.


Molecular Weight (observed) of ERI-1: 42 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, ERI-1 (h): 293T Lysate: sc-114320 or JAR cell lysate: sc-2276.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



ERI-1 (H-135): sc-66881. Western blot analysis of ERI-1 expression in non-transfected: sc-117752 (A) and human ERI-1 transfected: sc-114320 (B) 293T whole cell lysates.



ERI-1 (H-135): sc-66881. Western blot analysis of ERI-1 expression in NIH/3T3 whole cell lysate.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **ERI-1 (B-10): sc-137089** or **ERI-1 (C-8): sc-137099**, our highly recommended monoclonal alternatives to ERI-1 (H-135).