SANTA CRUZ BIOTECHNOLOGY, INC.

ESET (H-300): sc-66884



BACKGROUND

ERG-associated protein with SET domain (ESET), also designated Histone H3-K9 methyltransferase 4 or SET domain bifurcated 1, is a nuclear protein belonging to the histone-lysine methyltransferase family and to the Suvar3-9 subfamily. It is a highly conserved protein of 150 amino acids that has been implicated in chromatin structure modulation. ESET is excluded from cell nucleoli and areas of condensed chromatin and can associate with the nonpericentromeric regions of chromatin. The gene encoding for this protein, SETDB1, maps to chromosome 1q21.3. ESET is a histone methyltransferase, methylating Lys-9 of Histone H3 and mutations within the SETDB1 gene abolishes its methyltransferase activity. This methylation acts as a tag for epigenetic transcriptional repression by rounding up HP1 proteins to methylated histones. ESET is widely expressed with highest levels found in testis.

REFERENCES

- 1. Nomura, N., et al. 1994. Prediction of the coding sequences of unidentified human genes. II. The coding sequences of 40 new genes (KIAA0041-KIAA0080) deduced by analysis of cDNA clones from human cell line KG-1. DNA Res. 1: 223-229.
- 2. Harte, P.J., et al. 1999. Assignment of a novel bifurcated SET domain gene, SETDB1, to human chromosome band 1q21 by *in situ* hybridization and radiation hybrids. Cytogenet. Cell Genet. 84: 83-86.
- 3. Yang, L., et al. 2002. Molecular cloning of ESET, a novel histone H3specific methyltransferase that interacts with ERG transcription factor. Oncogene 21: 148-152.
- 4. Wang, H., et al. 2003. mAM facilitates conversion by ESET of dimethyl to trimethyl lysine 9 of histone H3 to cause transcriptional repression. Mol. Cell 12: 475-487.

CHROMOSOMAL LOCATION

Genetic locus: SETDB1 (human) mapping to 1q21.3; Setdb1 (mouse) mapping to 3 F2.1.

SOURCE

ESET (H-300) is a rabbit polyclonal antibody raised against amino acids 1-300 mapping at the N-terminus of ESET of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-66884 X, 200 $\mu\text{g}/0.1$ ml.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

ESET (H-300) is recommended for detection of ESET of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ESET (H-300) is also recommended for detection of ESET in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ESET siRNA (h): sc-45659, ESET siRNA (m): sc-45660, ESET shRNA Plasmid (h): sc-45669-SH, ESET shRNA Plasmid (m): sc-45660-SH, ESET shRNA (h) Lentiviral Particles: sc-45659-V and ESET shRNA (m) Lentiviral Particles: sc-45660-V.

ESET (H-300) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of ESET: 180 kDa.

Positive Controls: F9 cell lysate: sc-2245, MCF7 nuclear extract: sc-2149 or SW-13 cell lysate: sc-24778.

DATA





ESET (H-300): sc-66884. Western blot analysis of ESET expression in F9 whole cell lysate.

ESET (H-300): sc-66884. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization.

SELECT PRODUCT CITATIONS

- 1. Bilodeau, S., et al. 2009. SetDB1 contributes to repression of genes encoding developmental regulators and maintenance of ES cell state. Genes Dev. 23: 2484-2489.
- Jiang, Y., et al. 2010. Setdb1 histone methyltransferase regulates moodrelated behaviors and expression of the NMDA receptor subunit NR2B. J. Neurosci. 30: 7152-7167.
- Hong, W., et al. 2011. Epigenetic involvement of Alien/ESET complex in thyroid hormone-mediated repression of E2F1 gene expression and cell proliferation. Biochem. Biophys. Res. Commun. 415: 650-655.

MONOS Satisfation Guaranteed

Try **ESET (A-1):** sc-166621 or **ESET (G-4):** sc-271488, our highly recommended monoclonal aternatives to ESET (H-300).