

Mucin 13 (H-300): sc-66971

BACKGROUND

Mucins are epithelial glycoproteins with a high content of clustered oligosaccharides that are O-glycoside linked to tandem repeat peptides rich in threonine, serine and proline. Mucin 13 (MUC13), also designated down-regulated in colon cancer 1 (DRCC1), is an epithelial and hemopoietic type I membrane protein that undergoes secretion and influences gastrointestinal mucosa levels. It is most abundant in epithelial tissues of the gastrointestinal and respiratory tracts, such as large intestine and trachea, followed by kidney, small intestine, appendix and stomach. Mucin 13 is a good differentiation marker for gastrointestinal mucosa and may also indicate certain gastric tumors. It localizes to the apical membrane of both columnar and goblet cells in the gastrointestinal tract, and within goblet cell thecae. Mucin 13 is a cleaved protein, and the β subunit, containing the cytoplasmic tail, can homodimerize.

REFERENCES

- Williams, S.J., et al. 2001. Muc13, a novel human cell surface mucin expressed by epithelial and hemopoietic cells. *J. Biol. Chem.* 276: 18327-18336.
- Corrales, R.M., et al. 2003. Normal human conjunctival epithelium expresses MUC13, MUC15, MUC16 and MUC17 mucin genes. *Arch. Soc. Esp. Oftalmol.* 78: 375-381.
- Carraway, K.L., et al. 2003. Cell signaling through membrane mucins. *Bioessays* 25: 66-71.
- Packer, L.M., et al. 2004. Expression of the cell surface mucin gene family in adenocarcinomas. *Int. J. Oncol.* 25: 1119-1126.
- Byrd, J.C., et al. 2004. Mucins and mucin binding proteins in colorectal cancer. *Cancer Metastasis Rev.* 23: 77-99.
- Hollingsworth, M.A., et al. 2004. Mucins in cancer: protection and control of the cell surface. *Nat. Rev. Cancer* 4: 45-60.
- Shimamura, T., et al. 2005. Overexpression of MUC13 is associated with intestinal-type gastric cancer. *Cancer Sci.* 96: 265-273.

CHROMOSOMAL LOCATION

Genetic locus: MUC13 (human) mapping to 3q21.2.

SOURCE

Mucin 13 (H-300) is a rabbit polyclonal antibody raised against amino acids 19-318 mapping within an N-terminal extracellular domain of Mucin 13 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Mucin 13 (H-300) is recommended for detection of Mucin 13 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for Mucin 13 siRNA (h): sc-45690, Mucin 13 shRNA Plasmid (h): sc-45690-SH and Mucin 13 shRNA (h) Lentiviral Particles: sc-45690-V.

Molecular Weight of Mucin 13: 58 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



Mucin 13 (H-300): sc-66971. Immunoperoxidase staining of formalin fixed, paraffin-embedded human rectum tissue showing membrane and cytoplasmic staining of glandular cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Mucin 13 (D-5): sc-373857**, our highly recommended monoclonal alternative to Mucin 13 (H-300).