

Myosin Ia (H-90): sc-66979

BACKGROUND

Actin is a highly conserved protein that is expressed in all eukaryotic cells. Actin filaments can form both stable and labile structures and are crucial components of microvilli and the contractile apparatus of muscle cells. Troponin facilitates interaction between Actin and Myosin by binding to Ca^{2+} . Troponin is made up of at least two subunits, which are divergent in cardiac muscle, fast skeletal muscle and slow skeletal muscle. Myosin is a hexamer of two heavy chains (MHC) and four light chains (MLC) that interacts with Actin to generate the force for diverse cellular movements, including cytokinesis, phagocytosis and muscle contraction. Myosin Ia (MYO1A) is also designated Brush border myosin I or myosin I heavy chain. MYO1A, the gene encoding for the Myosin Ia protein, localizes to chromosome 12q13.3. Mutations in the MYO1A gene may cause autosomal dominant nonsyndromic sensorineural deafness (DFNA).

REFERENCES

- Hasson, T., Skowron, J.F., Gilbert, D.J., Avraham, K.B., Perry, W.L., Bement, W.M., Anderson, B.L., Sherr, E.H., Chen, Z.Y., Greene, L.A., Ward, D.C., Corey, D.P., Mooseker, M.S., Copeland, N.G. and Jenkins, N.A. 1996. Mapping of unconventional myosins in mouse and human. *Genomics* 36: 431-439.
- Li, W., Wang, J., Coluccio, L.M., Matsudaira, P. and Grand, R.J. 2000. Brush border myosin I (BBMI): a basally localized transcript in human jejunal enterocytes. *J. Histochem. Cytochem.* 48: 89-94.
- Hawkins, C.J., Silke, J., Verhagen, A.M., Foster, R., Ekert, P.G. and Ashley, D.M. 2001. Analysis of candidate antagonists of IAP-mediated caspase inhibition using yeast reconstituted with the mammalian Apaf-1-activated apoptosis mechanism. *Apoptosis* 6: 331-338.
- Donaudy, F., Ferrara, A., Esposito, L., Hertzano, R., Ben-David, O., Bell, R.E., Melchionda, S., Zelante, L., Avraham, K.B. and Gasparini, P. 2003. Multiple mutations of MYO1A, a cochlear-expressed gene, in sensorineural hearing loss. *Am. J. Hum. Genet.* 72: 1571-1577.
- SWISS-PROT/TrEMBL (Q9UBC5). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: MYO1A (human) mapping to 12q13.3; Myo1a (mouse) mapping to 10 D3.

SOURCE

Myosin Ia (H-90) is a rabbit polyclonal antibody raised against amino acids 761-850 mapping within an internal region of Myosin Ia of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Myosin Ia (H-90) is recommended for detection of Myosin Ia of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Myosin Ia siRNA (h): sc-44596, Myosin Ia siRNA (m): sc-44597, Myosin Ia shRNA Plasmid (h): sc-44596-SH, Myosin Ia shRNA Plasmid (m): sc-44597-SH, Myosin Ia shRNA (h) Lentiviral Particles: sc-44596-V and Myosin Ia shRNA (m) Lentiviral Particles: sc-44597-V.


Molecular Weight of Myosin Ia: 110 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Hep G2 cell lysate: sc-2227 or K-562 whole cell lysate: sc-2203.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Myosin Ia (H-90): sc-66979. Western blot analysis of Myosin Ia expression in HeLa (A), Hep G2 (B), Raji (C), K-562 (D), NCI-H929 (E) and Jurkat (F) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.