

Pirh2 (FL-261): sc-67033

BACKGROUND

Pirh2, also known as androgen receptor N-terminal-interacting protein (ARNIP), ZN363 or CHIMP, has p53-induced ubiquitin-protein ligase activity, promoting p53 degradation. The protein physically interacts with p53 and the resulting degradation of p53 renders Pirh2 an oncogenic protein, as the loss of p53 function contributes to malignant tumor development. The gene encoding for the protein maps to chromosome 4q21.1 and transcription of this gene is regulated by p53. Pirh2 expression decreases the level of p53, and a decrease of endogenous Pirh2 expression increases p53 levels. Pirh2 is therefore considered, together with MDM2, to act as a negative regulator of p53 function.

CHROMOSOMAL LOCATION

Genetic locus: RCHY1 (human) mapping to 4q21.1; Rchy1 (mouse) mapping to 5 E2.

SOURCE

Pirh2 (FL-261) is a rabbit polyclonal antibody raised against amino acids 1-261 representing full length Pirh2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-67033 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Pirh2 (FL-261) is recommended for detection of Pirh2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Pirh2 (FL-261) is also recommended for detection of Pirh2 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for Pirh2 siRNA (h): sc-45424, Pirh2 siRNA (m): sc-45425, Pirh2 shRNA Plasmid (h): sc-45424-SH, Pirh2 shRNA Plasmid (m): sc-45425-SH, Pirh2 shRNA (h) Lentiviral Particles: sc-45424-V and Pirh2 shRNA (m) Lentiviral Particles: sc-45425-V.

Pirh2 (FL-261) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Pirh2: 30 kDa.

Positive Controls: Pirh2 (m2): 293T Lysate: sc-122593.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



Pirh2 (FL-261): sc-67033. Western blot analysis of Pirh2 expression in non-transfected: sc-117752 (A) and mouse Pirh2 transfected: sc-122593 (B) 293T whole cell lysates.



Pirh2 (FL-261): sc-67033. Immunoperoxidase staining of formalin fixed, paraffin-embedded human adrenal gland tissue showing cytoplasmic staining of glandular cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Pirh2 (D-12): sc-374505** or **Pirh2 (E-11): sc-166901**, our highly recommended monoclonal alternatives to Pirh2 (FL-261).