robo4 (M-235): sc-67057



The Power to Question

BACKGROUND

Secreted leucine-rich repeat-containing proteins 1 through 3 (Slit1-3) are secreted glycoproteins that influence axonal guidance and mediate normal neural progression by acting as high-affinity signaling ligands for the repulsive guidance receptors, robo1 and robo2 (also designated roundabout 1 and 2). Interactions between the robo receptor and Slit ligand families of proteins initiate signaling cascades that repel axonal outgrowth. The arrangement of the extracellular domains of robo4 diverges significantly from that of all other robo family members. Robo4 is the only robo family member expressed in primary endothelial cells. It binds Slit and inhibits cellular migration in a heterologous expression system. Together, the robo proteins prescribe developmental paths during neural development.

REFERENCES

- 1. Huminiecki, L., et al. 2002. Magic roundabout is a new member of the roundabout receptor family that is endothelial specific and expressed at sites of active angiogenesis. Genomics 79: 547-552.
- 2. Park, K.W., et al. 2003. Robo4 is a vascular-specific receptor that inhibits endothelial migration. Dev. Biol. 261: 251-267.
- Middleton, R., et al. 2003. Improving the comparative map of porcine chromosome 9 with respect to human chromosomes 1, 7 and 11. Cytogenet. Genome Res. 102: 128-132.
- Suchting, S., et al. 2005. Soluble robo4 receptor inhibits in vivo angiogenesis and endothelial cell migration. FASEB J. 19: 121-123.
- 5. Seth, P., et al. 2005. Magic roundabout, a tumor endothelial marker: expression and signaling. Biochem. Biophys. Res. Commun. 332: 533-541.

CHROMOSOMAL LOCATION

Genetic locus: Robo4 (mouse) mapping to 9 A4.

SOURCE

robo4 (M-235) is a rabbit polyclonal antibody raised against amino acids 546-780 mapping within an internal region of robo4 of mouse origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

robo4 (M-235) is recommended for detection of robo4 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for robo4 siRNA (m): sc-44501, robo4 shRNA Plasmid (m): sc-44501-SH and robo4 shRNA (m) Lentiviral Particles: sc-44501-V.


Molecular Weight of robo4: 160 kDa.

Positive Controls: rat heart extract: sc-2393 or A-10 cell lysate: sc-3806.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



robo4 (M-235): sc-67057. Western blot analysis of robo4 expression in rat heart tissue extract.

SELECT PRODUCT CITATIONS

 Zhao, X.C., et al. 2013. Isoflurane post-conditioning protects primary cultures of cortical neurons against oxygen and glucose deprivation injury via upregulation of Slit2/Robo1. Brain Res. 1537: 283-289.



Try **robo4 (D-3): sc-166872**, our highly recommended monoclonal alternative to robo4 (M-235).

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com