

# Sall1 (H-65): sc-67087

## BACKGROUND

Sall1 (Sall1, sal-like 1, TBS, HSA1) and Sall2 (Sall2, sal-like 2, HSA2, p150 (Sal2)) are mammalian homologs of the *Drosophila* region-specific homeotic gene spalt (sal), which encodes a zinc finger-containing transcription regulator. *Drosophila* spalt (sal) is an essential genetic component required for the specification of posterior head and anterior tail as opposed to trunk. Mammalian Sall1 may mediate higher order chromatin structure and may be a component of a distinct heterochromatin-dependent silencing process. Sall1 is present in kidney, brain and liver. Sall2 is a p53-independent regulator of p21 and Bax, and can function in some cell types as a regulator of cell growth and survival. Human Sall2 is present in brain, heart, kidney or pancreas. Sall1 and Sall2 are expressed in different areas of the fetal brain that may represent distinct sets of neurons.

## REFERENCES

- Nielsen, T.O., et al. 2003. Tissue microarray validation of epidermal growth factor receptor and Sall2 in synovial sarcoma with comparison to tumors of similar histology. *Am. J. Pathol.* 163: 1449-1456.
- Sato, A., et al. 2003. Zinc finger protein Sall2 is not essential for embryonic and kidney development. *Mol. Cell. Biol.* 23: 62-69.
- Wabbels, B.K., et al. 2004. Clinical and molecular genetic findings in isolated sporadic Duane syndrome. *Klin. Monatsbl. Augenheilkd.* 221: 849-853.
- Wabbels, B.K., et al. 2004. No evidence of Sall4-mutations in isolated sporadic duane retraction "syndrome" (DURS). *Am. J. Med. Genet.* 131: 216-218.
- Borozdin, W., et al. 2004. Novel mutations in the gene Sall4 provide further evidence for acro-renal-ocular and Okihiro syndromes being allelic entities, and extend the phenotypic spectrum. *J. Med. Genet.* 41: e102.
- Borozdin, W., et al. 2004. Sall4 deletions are a common cause of Okihiro and acro-renal-ocular syndromes and confirm haploinsufficiency as the pathogenic mechanism. *J. Med. Genet.* 41: e113.
- Kohlhase, J., et al. 2004. Mutations in Sall4 in malformed father and daughter postulated previously due to reflect mutagenesis by thalidomide. *Birth Defects Res. A Clin. Mol. Teratol.* 70: 550-551.

## CHROMOSOMAL LOCATION

Genetic locus: SALL1 (human) mapping to 16q12.1; Sall1 (mouse) mapping to 8 C3.

## SOURCE

Sall1 (H-65) is a rabbit polyclonal antibody raised against amino acids 1260-1324 mapping at the C-terminus of Sall1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-67087 X, 200 µg/0.1 ml.

## APPLICATIONS

Sall1 (H-65) is recommended for detection of Sall1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Sall1 (H-65) is also recommended for detection of Sall1 in additional species, including canine and porcine.

Suitable for use as control antibody for Sall1 siRNA (h): sc-45620, Sall1 siRNA (m): sc-45621, Sall1 shRNA Plasmid (h): sc-45620-SH, Sall1 shRNA Plasmid (m): sc-45621-SH, Sall1 shRNA (h) Lentiviral Particles: sc-45620-V and Sall1 shRNA (m) Lentiviral Particles: sc-45621-V.


Sall1 (H-65) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Positive Controls: Sall1 (h): 293T Lysate: sc-369360 or Y79 cell lysate: sc-2240.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Sall1 (H-65): sc-67087. Western blot analysis of Sall1 expression in non-transfected 293T: sc-117752 (A), human Sall1 transfected 293T: sc-369360 (B) and Y79 (C) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.