

# NMT1 (H-55): sc-67172

## BACKGROUND

N-terminal myristoylation is a cotranslational lipid modification which is crucial for the targeting and function of many signaling proteins. The N-myristoyltransferases (NMT 1 and NMT 2), also known as glycylopeptide N-tetradecanoyltransferases, are cytoplasmic proteins that belong to the NMT family of proteins. The proteins in this family catalyze the addition of a myristoyl group to the N-terminal glycine residue of eukaryotic, fungal and viral proteins. They are primarily detected in heart, gut, kidney, liver and placenta.

## REFERENCES

1. McIlhinney, R.A., et al. 1994. Characterization of a polyhistidine-tagged form of human myristoyl-CoA: protein N-myristoyltransferase produced in *Escherichia coli*. *Eur. J. Biochem.* 222: 137-146.
2. Weston, S.A., et al. 1998. Crystal structure of the anti-fungal target N-myristoyl transferase. *Nat. Struct. Biol.* 5: 213-221.

## CHROMOSOMAL LOCATION

Genetic locus: NMT1 (human) mapping to 17q21.31; Nmt1 (mouse) mapping to 11 E1.

## SOURCE

NMT1 (H-55) is a rabbit polyclonal antibody raised against amino acids 1-55 mapping at the N-terminus of NMT1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

NMT1 (H-55) is recommended for detection of NMT1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NMT1 (H-55) is also recommended for detection of NMT1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for NMT1 siRNA (h): sc-61132, NMT1 siRNA (m): sc-61133, NMT1 shRNA Plasmid (h): sc-61132-SH, NMT1 shRNA Plasmid (m): sc-61133-SH, NMT1 shRNA (h) Lentiviral Particles: sc-61132-V and NMT1 shRNA (m) Lentiviral Particles: sc-61133-V.


Molecular Weight of NMT1: 66 kDa.

Positive Controls: NMT1 (h): 293 Lysate: sc-110896, NMT1 (m): 293T Lysate: sc-125714 or mouse pancreas extract: sc-364244.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



NMT1 (H-55): sc-67172. Western blot analysis of NMT1 expression in non-transfected: sc-110760 (A) and human NMT1 transfected: sc-110896 (B) 293 whole cell lysates.



NMT1 (H-55): sc-67172. Western blot analysis of NMT1 expression in non-transfected: sc-117752 (A) and mouse NMT1 transfected: sc-125714 (B) 293T whole cell lysates and mouse pancreas tissue extract (C).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **NMT1 (E-9): sc-393702** or **NMT1 (B-11): sc-393744**, our highly recommended monoclonal alternatives to NMT1 (H-55).