

TIG3 (H-70): sc-67174

BACKGROUND

Retinoids act through ligand-dependent transcription factors, retinoid X receptor (RXRs) and retinoic acid receptors (RARs). Tazarotene-induced gene proteins (TIGs), also designated retinoic acid receptor responder proteins or RAR-responsive proteins, can be membrane bound or secreted. They act as tumor suppressor genes in human cancers and are highly expressed in skin, hair follicles, endothelial cells as well as pancreas, spleen and intestine. TIGs have also been implicated as growth regulators that mediate the growth suppressive effects of retinoids, and they are activated by tazarotene. TIG1 is a single pass type II membrane protein activated by tazarotene and RAR proteins. It belongs to the protease inhibitor I47 (Latexin) family of proteins. TIG2 is a secreted protein that is mainly expressed in epidermis, hair follicles and endothelial cells. TIG2 is inhibited in psoriatic lesions and is activated by tazarotene in skin rafts and in epidermis of psoriatic lesions. TIG3 acts as a growth regulator, as it is important for mediating the growth suppressive effects of retinoids. This protein, which is widely expressed in most tissues (although not detected in heart, testis and brain), is activated by tazarotene and belongs to the H-rev107 family of proteins.

REFERENCES

- DiSepio, D., et al. 1998. Identification and characterization of a retinoid-induced class II tumor suppressor/growth regulatory gene. *Proc. Natl. Acad. Sci. USA* 95: 14811-14815.
- Tokumaru, Y., et al. 2004. Optimal use of a panel of methylation markers with GSTP1 hypermethylation in the diagnosis of prostate adenocarcinoma. *Clin. Cancer Res.* 10: 5518-5522.
- Youssef, E.M., et al. 2004. Hypermethylation and silencing of the putative tumor suppressor tazarotene-induced gene 1 in human cancers. *Cancer Res.* 64: 2411-2417.
- Takai, N., et al. 2005. Discovery of epigenetically masked tumor suppressor genes in endometrial cancer. *Mol. Cancer Res.* 3: 261-269.
- Aagaard, A., et al. 2005. An inflammatory role for the mammalian carboxypeptidase inhibitor Latexin: relationship to cystatins and the tumor suppressor TIG1. *Structure* 13: 309-317.
- Kwong, J., et al. 2005. Silencing of the retinoid response gene TIG1 by promoter hypermethylation in nasopharyngeal carcinoma. *Int. J. Cancer.* 113: 386-392.

CHROMOSOMAL LOCATION

Genetic locus: RARRES3 (human) mapping to 11q23.

SOURCE

TIG3 (H-70) is a rabbit polyclonal antibody raised against amino acids 42-111 mapping within an internal region of TIG3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

TIG3 (H-70) is recommended for detection of TIG3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for TIG3 siRNA (h): sc-61690, TIG3 shRNA Plasmid (h): sc-61690-SH and TIG3 shRNA (h) Lentiviral Particles: sc-61690-V.

Molecular Weight of TIG3: 18.2 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TIG3 (H-70): sc-67174. Western blot analysis of TIG3 expression in 293T (A) and K-562 (B) whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **TIG3 (LW-M10): sc-135579**, our highly recommended monoclonal alternative to TIG3 (H-70).