



LINE-1 (M-300): sc-67198

BACKGROUND

Mammalian LINE-1 (L1) elements belong to the superfamily of autonomously replicating retrotransposable elements that lack the long terminal repeated (LTR) sequences typical of retroviruses and retroviral-like retrotransposons. The non-LTR superfamily is ubiquitous in nature and constitutes 20% or more of some mammalian genomes and about 5% of the human genome. Therefore, LINE-1 elements presumably have had a profound, perhaps defining, effect on the evolution, structure and function of mammalian genomes. LINE-1 elements contain regulatory signals and encode two proteins: an RNA-binding protein and an integrase-replicase, as the retrovirus harbors both endonuclease and reverse transcriptase capabilities.

REFERENCES

- Dombroski, B.A., Mathias, S.L., Nanthakumar, E., Scott, A.F. and Kazazian, H.H. 1992. Isolation of an active human transposable element. *Science* 254: 1805-1808.
- Dombroski, B.A., Scott, A.F. and Kazazian, H.H. 1993. Two additional potential retrotransposons isolated from a human L1 subfamily that contains an active retrotransposable element. *Proc. Nat. Acad. Sci. USA* 90: 6513-6517.
- Brouha, B., Schustak, J., Badge, R.M., Lutz-Prigge, S., Farley, A.H., Moran, J.V. and Kazazian, H.H. 2003. Hot L1s account for the bulk of retrotransposition in the human population. *Proc. Nat. Acad. Sci. USA* 100: 5280-5285.
- Kulpa, D.A. and Moran, J.V. 2005. Ribonucleoprotein particle formation is necessary but not sufficient for LINE-1 retrotransposition. *Hum. Mol. Genet.* 14: 3237-3248.
- Roman-Gomez, J., Jimenez-Velasco, A., Agirre, X., Cervantes, F., Sanchez, J., Garate, L., Barrios, M., Castillejo, J.A., Navarro, G., Colomer, D., Prosper, F., Heiniger, A. and Torres, A. 2005. Promoter hypomethylation of the LINE-1 retrotransposable elements activates sense/antisense transcription and marks the progression of chronic myeloid leukemia. *Oncogene* 24: 7213-7223.
- Gilbert, N., Lutz, S., Morrish, T.A. and Moran, J.V. 2005. Multiple fates of L1 retrotransposition intermediates in cultured human cells. *Mol. Cell. Biol.* 25: 7780-7795.
- Han, J.S. and Boeke, J.D. 2005. LINE-1 retrotransposons: modulators of quantity and quality of mammalian gene expression? *Bioessays* 27: 775-784.
- Han, K., Sen, S.K., Wang, J., Callinan, P.A., Lee, J., Cordaux, R., Liang, P. and Batzer, M.A. 2005. Genomic rearrangements by LINE-1 insertion-mediated deletion in the human and chimpanzee lineages. *Nucleic Acids Res.* 33: 4040-4052.
- Grahn, R.A., Rinehart, T.A., Cantrell, M.A. and Wichman, H.A. 2005. Extinction of LINE-1 activity coincident with a major mammalian radiation in rodents. *Cytogenet. Genome Res.* 110: 407-415.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

LINE-1 (M-300) is a rabbit polyclonal antibody raised against amino acids 982-1281 mapping at the C-terminus of LINE-1 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

LINE-1 (M-300) is recommended for detection of LINE-1 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for LINE-1 siRNA (m): sc-60937.

Molecular Weight of LINE-1: 147 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.