

FOXN2 (E-15): sc-67487

BACKGROUND

The forkhead domain-containing gene family (Fox) comprises over 20 members in mammals and is defined by a conserved 110 amino-acid motif containing a winged helix structure DNA-binding domain. The members of this gene family have been implicated as key regulators of embryogenesis, cell cycling, cell lineage restriction and cancer. As such, FOXN2 contains a domain with homology to the forkhead DNA binding domain. FOXN2, or Human T-cell leukemia virus enhancer factor, is a 341 amino acid protein mapping to human gene FOXN2, which has been localized to human chromosome 2p16.3. This protein, encoded by a 1239-bp cDNA isolated from the Jurkat cDNA library, is capable of binding to a region of the human T-cell leukemia virus long terminal repeat (HTLV-I LTR) located between amino acids 155 and 117. This purine-rich region is important in the regulation of gene expression by the Ets family of transcription factors. FOXN2 is a unique cellular gene that may function in the transcriptional regulation of HTLV-I LTR.

REFERENCES

- Li, C., Luskis, A.J., Sparkes, R., Tran, S.M. and Gaynor, R. 1992. Characterization and chromosomal mapping of the gene encoding the cellular DNA binding protein HTLF. *Genomics* 13: 658-664.
- Schlake, T., Schorpp, M. and Boehm, T. 2000. Formation of regulator/target gene relationships during evolution. *Gene* 256: 29-34.
- Ariyama, Y., Mori, T., Shinomiya, T., Sakabe, T., Fukuda, Y., Kanamaru, A., Yamada, Y., Isobe, M., Seto, M., Nakamura, Y. and Inazawa, J. 2000. Chromosomal imbalances in adult T-cell leukemia revealed by comparative genomic hybridization: gains at 14q and 2p16-22 in cell lines. *J. Hum. Genet.* 44: 357-363.
- Boehm, T., Bleul, C.C. and Schorpp, M. 2003. Genetic dissection of thymus development in mouse and zebrafish. *Immunol. Rev.* 195: 15-27.
- Katoh, M. and Katoh, M. 2004. Characterization of human FOXN4 gene in silico. *Int. J. Mol. Med.* 14: 949-953.
- Katoh, M. and Katoh, M. 2004. Human FOX gene family (Review). *Int. J. Oncol.* 25: 1495-1500.
- Schuff, M., Rössner, A., Donow, C. and Knöchel, W. 2006. Temporal and spatial expression patterns of FOXN genes in *Xenopus laevis* embryos. *Int. J. Dev. Biol.* 50: 429-434.

CHROMOSOMAL LOCATION

Genetic locus: FOXN2 (human) mapping to 2p16.3.

SOURCE

FOXN2 (E-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of FOXN2 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-67487 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-67487 X, 200 µg/0.1 ml.

APPLICATIONS

FOXN2 (E-15) is recommended for detection of FOXN2 of human and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

FOXN2 (E-15) is also recommended for detection of FOXN2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for FOXN2 siRNA (h): sc-62341, FOXN2 shRNA Plasmid (h): sc-62341-SH and FOXN2 shRNA (h) Lentiviral Particles: sc-62341-V.

FOXN2 (E-15) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of FOXN2: 37 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.