

PIWIL4 (Y-18): sc-67593

BACKGROUND

PIWIL4 (PIWI-like protein 4), also known as HIWI2 is a 852 amino acid protein that belongs to the argonaute family. PIWIL4 contains one PAZ domain and one PIWI domain and is essential for the maintenance of germline stem cells. PIWIL4 is a cytoplasmic protein that is expressed in adult testis. It regulates spermatogenesis and primordial germ cell production and has an essential role in meiotic differentiation of spermatocytes and in self-renewal of spermatogonial stem cells. PIWIL4-null mice are of normal size and weight and have the expected life span. Homozygous PIWIL4-null females are fertile and have no obvious defects. However, PIWIL4-deficient males are infertile and show a meiotic progression defect in early prophase of meiosis I and progressive loss of germ cells with age. Mutant males show elevated expression of LINE-1 and intracisternal A particle (IAP) element transcripts in germ cell lineages. The gene encoding PIWIL4 maps to human chromosome 11.

CHROMOSOMAL LOCATION

Genetic locus: PIWIL4 (human) mapping to 11q21; Piwil4 (mouse) mapping to 9 A2.

SOURCE

PIWIL4 (Y-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PIWIL4 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-67593 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

PIWIL4 (Y-18) is recommended for detection of PIWIL4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PIWIL4 (Y-18) is also recommended for detection of PIWIL4 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PIWIL4 siRNA (h): sc-62458, PIWIL4 siRNA (m): sc-62459, PIWIL4 shRNA Plasmid (h): sc-62458-SH, PIWIL4 shRNA Plasmid (m): sc-62459-SH, PIWIL4 shRNA (h) Lentiviral Particles: sc-62458-V and PIWIL4 shRNA (m) Lentiviral Particles: sc-62459-V.


Molecular Weight of PIWIL4: 97 kDa.

Positive Controls: NTERA-2 cl.D1 whole cell lysate: sc-364181, WI-38 whole cell lysate: sc-364260 or PC-3 cell lysate: sc-2220.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PIWIL4 (Y-18): sc-67593. Western blot analysis of PIWIL4 expression in NTERA-2 cl.D1 (A), WI-38 (B) and PC-3 (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE** Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **PIWIL4 (10G9B11): sc-517215**, our highly recommended monoclonal alternative to PIWIL4 (Y-18).