

MLTK (K-17): sc-67684

BACKGROUND

MLTK (mitogen-activated protein kinase kinase kinase MLT, leucine zipper and sterile α motif-containing kinase, HCCS-4) is an 800 amino acid protein encoded by the human gene ZAK. MLTK belongs to the protein kinase superfamily, STE Ser/Thr protein kinase family, MAP kinase kinase kinase subfamily and contains one protein kinase domain and one SAM (sterile α motif) domain. MLTK is a cytoplasmic protein found as a homodimer. It will translocate to the nucleus upon ultraviolet B irradiation. It is a stress-activated component of a protein kinase signal transduction cascade and helps regulate the JNK and p38 pathways. MLTK also has a role in the regulation of S and G₂ cell cycle checkpoint by direct phosphorylation of Chk2. Isoform 1 causes cell shrinkage and disruption of Actin stress fibers and may have a role in neoplastic cell transformation and cancer development. Isoform 1 also phosphorylates Histone H3 at Ser 28.

CHROMOSOMAL LOCATION

Genetic locus: ZAK (human) mapping to 2q24.2; Zak (mouse) mapping to 2 C3.

SOURCE

MLTK (K-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MLTK of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-67684 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

MLTK (K-17) is recommended for detection of MLTK of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MLTK (K-17) is also recommended for detection of MLTK in additional species, including canine, bovine, porcine and avian.


Suitable for use as control antibody for MLTK siRNA (h): sc-62625, MLTK siRNA (m): sc-62626, MLTK shRNA Plasmid (h): sc-62625-SH, MLTK shRNA Plasmid (m): sc-62626-SH, MLTK shRNA (h) Lentiviral Particles: sc-62625-V and MLTK shRNA (m) Lentiviral Particles: sc-62626-V.

Molecular Weight of MLTK: 91 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MLTK (K-17): sc-67684. Western blot analysis of MLTK expression in NIH/3T3 whole cell lysate.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **MLTK (A-2): sc-390924**, our highly recommended monoclonal alternative to MLTK (K-17).