



Ste5 (yN-19): sc-6778

BACKGROUND

MAP kinase cascades, consisting of a mitogen-activated protein kinase (also called ERK, for extracellular regulated kinase) and one or more upstream regulatory kinases (MAPKKs), play an integral role in signal transduction. The best characterized MAP kinase pathway to date is the yeast pheromone response pathway. Extracellular pheromones bind to the receptors Ste2 and Ste3 on the cell surface. Stimulation of these receptors leads to the activation of the kinase Ste20, which leads to the activation of the MAPKK Ste11. Upon phosphorylation, Ste11 activates the MAPKK Ste7, which in turn activates the MAP kinases Fus3 (also called Dac2) and Kss1. These MAP kinases activate the transcription factor Ste12, which upregulates mating-specific genes, and Far1, which arrests the cell cycle. Cross-talk between this and other MAP kinase cascades is restricted by Ste5 (also called Nul3), which is thought to bind these components together in a functional complex.

REFERENCES

- Errede, B. and Ammerer, G. 1989. Ste12, a protein involved in cell-type-specific transcription and signal transduction in yeast, is part of protein-DNA complexes. *Genes Dev.* 3: 1349-1361.
- Leberer, E., Dignard, D., Marcus, D., Thomas, D.Y. and Whiteway, M. 1992. The protein kinase homologue Ste20p is required to link the yeast pheromone response G protein $\beta\gamma$ subunits to downstream signalling components. *EMBO J.* 11: 4815-4824.
- Peter, M., Gartner, A., Horecka, J., Ammerer, G. and Herskowitz, I. 1993. Far1 links the signal transduction pathway to the cell cycle machinery in yeast. *Cell* 73: 747-760.
- Printen, J.A. and Sprague, G.F., Jr. 1994. Protein-protein interactions in the yeast pheromone response pathway: Ste5p interacts with all members of the MAP kinase cascade. *Genetics* 138: 609-619.
- Ferguson, B., Horecka, J., Printen, J., Schultz, J., Stevenson, B.J. and Sprague, G.F., Jr. 1994. The yeast pheromone response pathway: new insights into signal transmission. *Cell. Mol. Biol. Res.* 40: 223-228.
- Wu, C., Whiteway, M., Thomas, D.Y. and Leberer, E. 1995. Molecular characterization of Ste20p, a potential mitogen-activated protein of extracellular signal-regulated kinase kinase (MEK) kinase kinase from *Saccharomyces cerevisiae*. *J. Biol. Chem.* 270: 15984-15992.
- Bardwell, L., Cook, J.G., Chang, E.C., Cairns, B.R. and Thorner, J. 1996. Signaling in the yeast pheromone response pathway: specific and high-affinity interaction of the mitogen-activated protein (MAP) kinases Kss1 and Fus3 with the upstream MAP kinase kinase Ste7. *Mol. Cell. Biol.* 16: 3637-3650.

SOURCE

Ste5 (yN-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Ste5 of *Saccharomyces cerevisiae* origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6778 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Ste5 (yN-19) is recommended for detection of Ste5 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.