# Ste18 (yC-19): sc-6787



The Power to Question

#### **BACKGROUND**

MAP kinase cascades, consisting of a mitogen-activated protein kinase (also called ERK, for extracellular regulated kinase) and one or more upstream regulatory kinases (MAPKKs), play an integral role in signal transduction. One of the best characterized MAP kinase pathways is the yeast pheromone response pathway. Extracellular pheromones bind to the receptors Ste2 and Ste3, which couple to a heterotrimeric G protein. The G protein consists of  $\alpha$ ,  $\beta$  and  $\gamma$  subunits, respectively designated Gpa1 (also called Scg1, Cdc70 or Dac1), Ste4 and Ste18. Activation of this G protein leads to the activation of the MAPKKK Ste11, which then activates the MAPKK Ste7. Activated Ste7 phosphorylates the MAP kinases Fus3 (also called Dac2) and Kss1. These MAP kinases activate the transcription factor Ste12, which upregulates mating-specific genes and Far1, which is involved in cell cycle arrest.

## **REFERENCES**

- Nakayama, N., Mayajima, A. and Arai, K. 1985. Nucleotide sequences of STE2 and STE3, cell type-specific sterile genes from *Saccharomyces* cerevisiae. EMBO J. 4: 2643-2648.
- 2. Dietzel, C. and Kurjan, J. 1987. The yeast SCG1 gene: a G  $\alpha$ -like protein implicated in the  $\alpha$  and  $\alpha$ -factor response pathway. Cell 50: 1001-1010.
- 3. Whiteway, M., Hougan, L., Dignard, D., Thomas, D.Y., Bell, L., Saari, G.C., Grant, F.J., O'Hara, P. and MacKay, V.L. 1989. The STE4 and STE18 genes of yeast encode potential  $\beta$  and  $\gamma$  subunits of the mating factor receptor-coupled G protein. Cell 56: 467-477.
- Dolan, J.W., Kirkman, C. and Fields, S. 1989. The yeast Ste12 protein binds to the DNA sequence mediating pheromone induction. Proc. Natl. Acad. Sci. USA 86: 5703-5707.
- Elion, E.A., Grisafi, P.L. and Fink, G.R. 1990. Fus3 encodes a Cdc2+/Cdc28related kinase required for the transition from mitosis into conjugation. Cell 60: 649-664.
- Peter, M., Gartner, A., Horecka, J., Ammerer, G. and Herskowitz, I. 1993.
  Far1 links the signal transduction pathway to the cell cycle machinery in yeast. Cell 73: 747-760.
- 7. Ferguson, B., Horecka, J., Printen, J., Schultz, J., Stevenson, B.J. and Sprague, G.F., Jr. 1994. The yeast pheromone response pathway: new insights into signal transmission. Cell. Mol. Biol. Res. 40: 223-228.
- Bardwell, L., Cook, J.G., Chang, E.C., Cairns, B.R. and Thorner, J. 1996. Signaling in the yeast pheromone response pathway: specific and highaffinity interaction of the mitogen-activated protein (MAP) kinases Kss1 and Fus3 with the upstream MAP kinase kinase Ste7. Mol. Cell Biol. 16: 3637-3650.

## **SOURCE**

Ste18 (yC-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Ste18 of *Saccharomyces cerevisiae* origin.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **PRODUCT**

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6787 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## **APPLICATIONS**

Ste18 (yC-19) is recommended for detection of Ste18 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

## **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com