

CYB5R3 (D-15): sc-68062

BACKGROUND

CYB5R3 (NADH-cytochrome b5, reductase 3, diaphorase 1) is a 301 amino acid protein encoded by the human gene CYB5R3. CYB5R3 belongs to the flavoprotein pyridine nucleotide cytochrome reductase family and has two naturally occurring isoforms. Isoform 1 is anchored to the cytoplasmic side of the endoplasmic reticulum membrane and mitochondrion outer membrane, while isoform 2 is the soluble form found in erythrocytes. CYB5R3 is involved in the desaturation and elongation of fatty acids, cholesterol biosynthesis, drug metabolism and, in erythrocytes, methemoglobin reduction. A serine residue at position 117 seems to only be found in persons of African origin. The allele frequency is 0.23 in African Americans. It is not found in Caucasians, Asians, Indo-Aryans or Arabs. This difference seems to have no effect on the enzyme activity. Defects in CYB5R3 are the cause of hereditary methemoglobinemia (HM). There are three forms of this disease: type 1 (HM1), in which the enzyme is only deficient in erythrocytes with a mild cyanosis; type 2 (HM2), in which the enzyme is completely deficient; and type 3 (HM3), where the deficiency is seen in all blood cells. Type 2 is a severe form accompanied by mental retardation and neurological impairment.

REFERENCES

- Farahani, P., Chiu, S., Bowlus, C.L., Boffelli, D., Lee, E., Fisler, J.S., Krauss, R.M. and Warden, C.H. 2004. Obesity in BSB mice is correlated with expression of genes for iron homeostasis and leptin. *Obes. Res.* 12: 191-204.
- Roma, G.W., Crowley, L.J., Davis, C.A. and Barber, M.J. 2005. Mutagenesis of Glycine 179 modulates both catalytic efficiency and reduced pyridine nucleotide specificity in cytochrome b5 reductase. *Biochemistry* 44: 13467-13476.
- Tonegawa, Y., Umeda, N., Hayakawa, T. and Ishibashi, T. 2005. Evaluation of data in terms of two-dimensional random walk model: interaction between NADH-cytochrome b5 reductase and cytochrome b5. *Biomed. Res.* 26: 207-212.
- Percy, M.J., Crowley, L.J., Davis, C.A., McMullin, M.F., Savage, G., Hughes, J., McMahon, C., Quinn, R.J., Smith, O., Barber, M.J. and Lappin, T.R. 2005. Recessive congenital methaemoglobinemia: functional characterization of the novel D239G mutation in the NADH-binding lobe of cytochrome b5 reductase. *Br. J. Haematol.* 129: 847-853.
- Percy, M.J., Crowley, L.J., Boudreaux, J. and Barber, M.J. 2006. Expression of a novel P275L variant of NADH:cytochrome b5 reductase gives functional insight into the conserved motif important for pyridine nucleotide binding. *Arch. Biochem. Biophys.* 447: 59-67.
- Roma, G.W., Crowley, L.J. and Barber, M.J. 2006. Expression and characterization of a functional canine variant of cytochrome b5 reductase. *Arch. Biochem. Biophys.* 452: 69-82.

CHROMOSOMAL LOCATION

Genetic locus: CYB5R3 (human) mapping to 22q13.2; Cyb5r3 (mouse) mapping to 15 E1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

CYB5R3 (D-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CYB5R3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-68062 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CYB5R3 (D-15) is recommended for detection of CYB5R3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CYB5R3 (D-15) is also recommended for detection of CYB5R3 in additional species, including canine.

Suitable for use as control antibody for CYB5R3 siRNA (h): sc-62173, CYB5R3 siRNA (m): sc-62174, CYB5R3 shRNA Plasmid (h): sc-62173-SH, CYB5R3 shRNA Plasmid (m): sc-62174-SH, CYB5R3 shRNA (h) Lentiviral Particles: sc-62173-V and CYB5R3 shRNA (m) Lentiviral Particles: sc-62174-V.

Molecular Weight of CYB5R3 isoforms: 32/34 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **CYB5R3 (G-11): sc-398043**, our highly recommended monoclonal alternative to CYB5R3 (D-15).