

# GIPC3 (V-18)-R: sc-68086-R

## BACKGROUND

The eukaryotic PDZ domain is a multifunctional protein-protein interacting motif that is found in a variety of proteins and is involved in both the clustering of signaling molecules and the organization of protein networks. GIPC3 (GIPC PDZ domain containing family, member 3) is a 312 amino acid protein that contains one PDZ domain and is a member of the GIPC family. Widely expressed with highest expression in small intestine and fetal spleen, GIPC3 may participate in signaling events throughout the cell via its central PDZ domain. Expression of GIPC3 is upregulated in melanoma, cervical, chronic myelogenous and gastric cancer cell lines, suggesting a possible role in carcinogenesis.

## REFERENCES

1. Ranganathan, R. and Ross, E.M. 1997. PDZ domain proteins: scaffolds for signaling complexes. *Curr. Biol.* 7: R770-R773.
2. Ponting, C.P., Phillips, C., Davies, K.E. and Blake, D.J. 1997. PDZ domains: targeting signalling molecules to sub-membranous sites. *Bioessays* 19: 469-479.
3. Saitoh, T., Mine, T. and Katoh, M. 2002. Molecular cloning and characterization of human GIPC3, a novel gene homologous to human GIPC1 and GIPC2. *Int. J. Oncol.* 20: 577-582.
4. Kirikoshi, H. and Katoh, M. 2002. Up-regulation of GIPC2 in human gastric cancer. *Int. J. Oncol.* 20: 1183-1187.
5. Kirikoshi, H. and Katoh, M. 2002. Expression of WNT7A in human normal tissues and cancer, and regulation of WNT7A and WNT7B in human cancer. *Int. J. Oncol.* 21: 895-900.
6. Saitoh, T., Mine, T. and Katoh, M. 2002. Molecular cloning and characterization of mouse Gipc3. *Int. J. Mol. Med.* 9: 251-256.
7. Kirikoshi, H. and Katoh, M. 2002. Expression of human GIPC1 in normal tissues, cancer cell lines, and primary tumors. *Int. J. Mol. Med.* 9: 509-513.
8. Katoh, M. 2002. GIPC gene family. *Int. J. Mol. Med.* 9: 585-589.

## CHROMOSOMAL LOCATION

Genetic locus: GIPC3 (human) mapping to 19p13.3; Gipc3 (mouse) mapping to 10 C1.

## SOURCE

GIPC3 (V-18)-R is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of GIPC3 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-68086 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

GIPC3 (V-18)-R is recommended for detection of GIPC3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GIPC3 (V-18)-R is also recommended for detection of GIPC3 in additional species, including canine and bovine.

Suitable for use as control antibody for GIPC3 siRNA (h): sc-62376, GIPC3 siRNA (m): sc-62377, GIPC3 shRNA Plasmid (h): sc-62376-SH, GIPC3 shRNA Plasmid (m): sc-62377-SH, GIPC3 shRNA (h) Lentiviral Particles: sc-62376-V and GIPC3 shRNA (m) Lentiviral Particles: sc-62377-V.

Molecular Weight of GIPC3: 34 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **GIPC3 (8H8): sc-517166**, our highly recommended monoclonal alternative to GIPC3 (V-18).