



Mkk2 (yC-17): sc-6811

BACKGROUND

Yeasts maintain the integrity of their cell walls via a MAP kinase cascade. This cascade consists of a MAP kinase (mitogen-activated protein kinase, also called ERK, for extracellular regulated kinase) as well as several upstream regulatory kinases (MAPKKs or MEKs, for MAP/ERK kinase). Pkc1 (also designated Sst1), a yeast homolog of the mammalian PKC α , β , and γ isoforms, transmits extracellular signals to Bck1, a MAPKKK (also called Slk1, Ssp31 or Las3). Bck1 then activates two MAPKKs, Mkk1 and Mkk2 (also referred to as Ssp32 and Ssp33, respectively). These in turn activate the MAP kinase Mpk1 (also called Slt2). Mutants lacking any component of this cascade exhibit a defect in cell lysis resulting from deficient cell wall synthesis. Bck2 (also designated Ctr7) has been identified as a suppressor of Pkc1 and Mpk1 deletions.

REFERENCES

- Lee, K.S., Irie, K., Gotoh, Y., Watanabe, Y., Araki, H., Eisuke, N., Matsumoto, K. and Levin, D.E. 1993. A yeast mitogen-activated protein kinase homolog (Mpk1p) mediates signalling by protein kinase C. *Mol. Cell. Biol.* 13: 3067-3075.
- Irie, K., Takase, M., Lee, K.S., Levin, D.E., Araki, H., Matsumoto, K. and Oshima, Y. 1993. MKK1 and MKK2, which encode *Saccharomyces cerevisiae* mitogen-activated protein kinase-kinase homologs, function in the pathway mediated by protein kinase C. *Mol. Cell. Biol.* 13: 3076-3083.
- Wu, J., Harrison, J.K., Dent, P., Lynch, K.R., Weber, M.J. and Sturgill, T.W. 1993. Identification and characterization of a new mammalian mitogen-activated protein kinase kinase, MKK2. *Mol. Cell. Biol.* 13: 4539-4548.
- Lee, K.S., Hines, L.K. and Levin, D.E. 1993. A pair of functionally redundant yeast genes (PPZ1 and PPZ2) encoding type 1-related protein phosphatases function within the PKC1-mediated pathway. *Mol. Cell. Biol.* 13: 5843-5853.
- Levin, D.E., Bowers, B., Chen, C.-Y., Kamada, Y. and Watanabe, M. 1994. Dissecting the protein kinase c/map kinase signalling pathway of *Saccharomyces cerevisiae*. *Cell. Mol. Biol. Res.* 40: 229-239.
- Watanabe, M., Chen, C.-Y. and Levin, D.E. 1994. *Saccharomyces cerevisiae* PKC1 encodes a protein kinase C (PKC) homolog with a substrate specificity similar to that of mammalian PKC. *J. Biol. Chem.* 269: 16829-16836.
- Soler, M., Plovins, A., Martin, H., Molina, M. and Nombela, C. 1995. Characterization of domains in the yeast MAP kinase Slt2 (Mpk1) required for functional activity and *in vivo* interaction with protein kinases Mkk1 and Mkk2. *Mol. Microbiol.* 17: 833-842.
- Paravicini, G. and Friedli, L. 1996. Protein-protein interactions in the yeast PKC1 pathway: Pkc1p interacts with a component of the MAP kinase cascade. *Mol. Gen. Genet.* 251: 682-691.
- Zarrov, P., Mazzoni, C. and Mann, C. 1996. The SLT2(MPK1) MAP kinase is activated during periods of polarized cell growth in yeast. *EMBO J.* 15: 83-91.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

Mkk2 (yC-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Mkk2 of *Saccharomyces cerevisiae* origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6811 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Mkk2 (yC-17) is recommended for detection of Mkk2 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.